

PREPRINT

Author-formatted, not peer-reviewed document posted on 01/08/2023

DOI: <https://doi.org/10.3897/arphapreprints.e110124>

Managing Invasive Wild Boars in Southern Brazil's Protected Areas: Challenges and Strategies

 **Matheus Etges, Demétrio L. Guadagnin, Andreas Kindel**

Managing Invasive Wild Boars in Southern Brazil's Protected Areas: Challenges and Strategies

Matheus Fragoso Etges^{1,2}, Demétrio Luis Guadagnin², Andreas Kindel³

¹ Corresponding author: matheus.etges@gmail.com.br; <https://orcid.org/0000-0001-5040-0158>

² Laboratório de Manejo e Conservação da Vida Silvestre. Department of Ecology, Federal University of Rio Grande do Sul, Av. Bento Gonçalves 9500, CEP 91501-970, Porto Alegre, RS, Brazil.

³ Núcleo de Ecologia de Rodovias e Ferrovias (NERF/UFRGS), Department of Ecology, Federal University of Rio Grande do Sul, Av. Bento Gonçalves 9500, CEP 91501-970, Porto Alegre, RS, Brazil.

Abstract

Invasive species pose significant threats to ecosystems and biodiversity, necessitating effective management strategies to mitigate their impacts. One such invasive species of concern is the wild boar in Brazil, which has the potential to cause widespread environmental changes. A national plan for monitoring and controlling invasive species, including the wild boar, was developed in response to this threat. Despite this initiative, uncertainties persist regarding the presence of wild boars in protected areas (PAs) and the effectiveness of current management actions. This study aimed to diagnose the situation of wild boars in protected areas within the southern region of Brazil, specifically focusing on their distribution, management techniques employed, and reasons for the lack of management action. An online questionnaire was sent to 297 PAs, with 134 responding. The findings revealed that wild boars were present in 39 of the surveyed PAs, but management efforts were only being carried out in 14 of them. Cages and corrals were identified as the most commonly used techniques, with corn serving as the preferred bait. The study identified two primary reasons for the lack of management action: the wild boar's low invasion intensity and management capacity limitations. To address these challenges effectively, the study advocates for a centralized organization of management actions and emphasizes the development of materials and resources to support successful management strategies. Implementing these measures is essential to safeguarding the conservation of ecosystems and vulnerable species in Brazil's protected areas and ensuring the long-term resilience of these valuable ecological assets.

Keywords: *Sus scrofa*, Online questionnaire, Invasive species, Conservation efforts, Management techniques, Wild Pigs

Introduction

Invasive alien species (IAS) are one of the most important direct drivers of biodiversity loss and ecosystem service change (Pyšek et al. 2020; Jaureguiberry et al. 2022), globally increasing at an unprecedented pace (Butchart et al. 2010). The strong links between invasions and other major drivers of change, such as global warming, pollution, overexploitation of resources, and habitat loss, are expected to increase the challenges (Simberloff et al. 2013; Spear et al. 2013), as well as the opportunities to manage (Dudley and Stolton 2010).

Risks associated with biological invasions tend to be greater in Protected Areas PAs both because of their importance for biodiversity conservation and because, once created, habitat loss, overexploitation, and pollution tend to be reduced (Klinger et al.

2006; Foxcroft et al. 2017; Liu et al. 2020; Ren et al. 2021). Risks even are greater if the invasive species has the potential to cause various effects on the landscape, such as wild boar (Barrios-Garcia and Ballari 2012; Risch et al. 2021).

The wild boar (*Sus scrofa*), one of the world's most widely distributed invasive exotic species (Long 2005), has a long history of invasion in Brazil's southern region (Hegel et al. 2022; La Sala et al. 2023). Wild pigs' fast expansion in Brazil was documented in the late 1980s - early 1990s, fueled by dispersion from neighboring countries and the introduction of leisure hunting and commercial interests (Deberdt and Scherer 2007; Salvador 2012; Hegel et al. 2022). Wild boar found a favorable breeding ground in the region, with a mosaic of agriculture and native forests supplying resources, and they still occupy these areas today (Hegel et al. 2019). These areas are also climatically similar to the ones the wild pigs are already adapted to (Sales et al. 2017; La Sala et al. 2023).

Controlling wild boars in PAs is challenged by constraints on techniques, resources, and management priorities. Although the recurrent calls for the importance of managing invasive species more efficiently in PAs and ideas to overcome management obstacles (Laurance et al. 2012; Tu and Robison 2013; Pyšek et al. 2020). Although there is a need to avoid invasive species' effects on conservation targets, assessing such effects is not always easy or possible (de Souza and Alves 2014; Keuling et al. 2016; Castilho et al. 2018). Managing invasive species is still infrequent (Kiringe et al. 2007; Genovesi and Monaco 2013). Andreu et al. (2009) listed eleven main obstacles to the efficient management of invasive plants in PAs, and they possibly apply also to the management of invasive animals: (i) the lack of capacity for mainstreaming IAS management into overall PA management, (ii) the limited capacity of staff at the site level, (iii) the low level of awareness, (iv) the gaps in information on IAS available to PA managers, (v) the lack of funding, (vii) legal or institutional impediments, (viii) and the clashes of interests between stakeholders. Despite the wild boar pervasive presence in protected areas throughout South America and Brazil (Sampaio and Schmidt 2013; La Sala et al. 2023), little is known about control efforts and management barriers in those PA systems. Worldwide the main techniques used for wild boar control can be divided into two major categories: lethal or non-lethal (West et al. 2009; Rosa et al. 2018; Gürtler et al. 2018; VerCauteren et al. 2019; Jori et al. 2021). Lethal techniques reduce abundance by increasing mortality (e.g., hunting, poisoning, trapping), while non-lethal techniques restrict resource access through movement restrictions or reduce fertility (Fryxell et al. 2014). In Brazil, it is legally allowed to control wild boars by hunting, either with or without the aid of dogs and live trapping. In this scenario, hunting is the primary technique used for controlling wild pigs in rural properties, while in PAs, trapping is routinely used (Rosa et al. 2018).

Based on documented records and questionnaires sent to managers, we describe the status of wild boar management in Protected Areas of southern Brazil and address the reasons behind the decision to manage or not to manage wild boards in protected areas. Due to the long history of the introduction and spread of wild boards in South Brazil, we consider that its presence inside the protected areas can be explained by habitat and landscape factors and that the perceived impact, in relation to other management demands, explains management prioritization. We expect that the management actions will differ between the PAs, where despite the techniques potentially being the same, the equipment, routine, and effort will be different. As possible justifications for non-

management, we expect to find answers similar to those found by Andreu et al. (2009) in the management of invasive plants: (i) the lack of capacity for mainstreaming IAS management into overall PA management, (ii) the limited capacity of staff at the site level, (iii) the low level of awareness, (iv) the gaps in information on IAS available to PA managers, (v) the lack of funding, (vi) legal or institutional impediments, (vii) and the clashes of interests between stakeholders.

Methods

Study area

The southern region of Brazil comprises the states of Rio Grande do Sul (RS), Santa Catarina (SC), and Paraná (PR), summing an area of 576,774 km². The region includes two biomes (Dinerstein et al. 2017): Tropical and Subtropical Grasslands, Savannas and Shrublands (Grasslands); and Tropical and Subtropical Moist Broadleaf Forests (Forests). They present approximately 3% and 10.3% of their coverage within protected areas, respectively (*Cadastro Nacional de Unidades de Conservação*).

The Brazilian system of protected areas includes federal, state, municipal, and private PAs, grouped into two major categories: Strictly Protected Areas and Sustainable Use Areas (Rylands and Brandon 2005), equivalent, respectively, to categories I to IV and categories V to VI of the International Union for Conservation of Nature (IUCN data).

Data collection

We compiled the list of protected areas in South Brazil from the National Catalog of Protected Areas (*Cadastro Nacional de Unidades de Conservação*) maintained by the Chico Mendes Institute of Biodiversity Conservation (<https://www.gov.br/mma/pt-br/assuntos/areasprotegidasecoturismo/plataforma-cnuc-1>) together with the information found in the state secretariats. We included in the study all areas: Federal, State and Municipal, and private or public. Marine conservation areas were excluded from the work. In all, we compiled 708 protected areas for this region.

We initially contacted the protected areas through e-mails available on the relevant authorities' website. For those without information, tried phone contacts and looked for and tried other e-mails and contacts searched on the web. For e-mails found, we send an online survey addressed to the PA managers. Four contact attempts were made with each PA (October and November 2019 and February and March 2020).

We structured the questionnaire in three sections on the Research Electronic Data Capture (REDCap) platform (Table 1; Supplementary Material). The first section encompassed questions about wild boar occurrence and its effects within the PA. The second section was dedicated to obtaining information about control techniques or reasons for not adopting control. Finally, the third section looked at the frequency of control campaigns, the number of animals managed, and their destination. The questionnaire primarily consists of multiple-choice questions and short texts (Table 1). However, in the first session, there are also two questions to understand the manager's perception of the effects of the wild boar compared to other generators of negative effects within the protected area. In these questions, the manager must choose a number from 0% to 100%, where the higher the number, the greater the responsibility of the wild boar in generating the negative effects.

We summarized answers using descriptive statistics. We grouped and named the answers to the open-ended questions a posteriori according to their similarity using the technique for qualitative analysis of Discourse Analysis, where the main ideas are extracted from the text to summarize and group responses into categories (Taylor 2013; Georgakopoulou 2019).

Table 1. Summary of questions asked in the survey. "Type" represents the types of responses accepted: "Text" correspond to short, typed responses accepting letters and numbers; "0-100" scale of numerical answers where 0 represents "Does not generate negative effects" and 100 represents "Greater driver of negative effects"; "Long Text" for answers where it is allowed to write up to 250 words; "Multiple Choice" when it is allowed to mark at least one of presented options. The "Others" option means that the option is not listed above, and typically, the next question asking for details will appear when checked.

First section questions/options	Type
- Are there wild boars in this area?	Yes/No
- What year was the first sighting?	Text
- Among all the factors causing adverse effects within the PA, how much is the wild boar responsible for?	0-100
- Of all the invasive species within the PA, how much does the wild boar cause adverse effects?	0-100
Second section questions/options	Type
- In the last year, was any management action taken to control the wild boar?	Yes/No
- Explain the reasons why control actions were not carried out.	Long Text
- Is there any specialized technical support that helps in the execution of the management project?	Yes/No
- This technical support is provided by:	Multiple Choice
University/ Government Agencies/ NGOs/ Companies/ Others	
- What types of actions were taken?	Multiple Choice
Hunting with dogs/ Hunting without dogs/ Stands/ Corral/ Cage/ Fencing/ Others	
- How many cages were used?	Text
- Do the cages used have different sizes?	Yes/No
- How many corrals were used?	Text
- Are the corrals different sizes?	Yes/No
- Where did the traps come from?	Multiple Choice
Store/ NGO/ Own Construction/ Others	
- Are baits used?	Yes/No
- What bait was used?	Text
- Where is the bait placed?	
At the entrance/ Around/ At the bottom/ Others	Multiple Choice
- When is the bait placed?	Text
E.g., Last day, There is always bait on site.	
Third section questions/options	Type

- What was the frequency of management campaigns?	Text
E.g., Once a month, every month, once a year.	
- What is the duration of the campaigns?	Text
E.g., Five days; One month.	
- How many people participate on average in the campaigns?	Text
- Describe the control procedure.	Long Text
- Is a record made of how many animals were captured?	Yes/No
- How many individuals were captured?	Text
- Is there any sanitary control or disease monitoring of captured individuals?	Yes/No
- What is the destination of the carcasses?	Text
- Space to be filled in if you want to report something or add some information.	Long Text

Results

We compiled a list of 701 protected areas from the National Catalog, of which nine were excluded because they were maritime (Table 2). From the final list, 297 protected areas were contacted, of which we got responses from 134 (Table 2). Concerning the public sphere, we were able to represent all management spheres of interest and all ecoregions/biogeographical provinces (Table 3). Due to only two responses for the private sphere (0.6%), these were not considered in this work (Table 2-3). In all cases, the respondents of the surveys were the protected area managers, and only finished questionnaires were considered in the results.

Table 2. Summary of search for PAs contact. Summaries of the contact process, questionnaire return, and inclusion of PAs. Total, represents the number of PAs registered for the administrative sphere, category of use and biomes. Excluded represents those that were not fetched for contact. Contacted represents those that it was possible to get a contact email. Unanswered and included represents the number of responses to the questionnaire. Inc./Total represents the percentage of returns in relation to the total recorded. Forest = Tropical and Subtropical Moist Broadleaf Forests; Grasslands = Tropical and Subtropical Grasslands, Savannas and Shrublands.

		Total	Excluded	Contacted	No-answer	Included	Inc./Total
Sphere	Federal	40	8	30	1	29	72.5%
	State	105	0	60	15	45	42.9%
	Municipal	220	1	164	106	58	26.4%
	Private	336	0	43	41	2	0.6%
Category	Strictly Protected	249	6	181	85	96	38.6%
	Sustainable Use	452	3	116	78	38	8.4%
Biome	Marine	9	9	-	-	-	-
	Forests	658	0	271	156	115	17.5%
	Grasslands	34	0	26	7	19	55.9%

Table 3. Summary of PAs characteristics used in this work. Number of protected areas included in the analyses and its representative percentage in relation to the registered total by category grouping. Forest = Tropical and Subtropical Moist Broadleaf Forests; Grasslands = Tropical and Subtropical Grasslands, Savannas and Shrublands. * Excluding private areas from the total and responses.

		Category		Biome	
		Strictly Protected	Sustainable Use	Forests	Grasslands
Sphere	Federal	16 (72.7%)	13 (72.2%)	25 (89.3%)	4 (100%)
	State	40 (52.6%)	5 (17.2%)	36 (37.9%)	9 (90%)
	Municipal	40 (26.5%)	18 (26.1%)	53 (25.2%)	8 (88.9%)
	Private	0 (0%)	2 (0.6%)	1 (0.3%)	1 (9.1%)
Biome	Forests	83 (36.9%)	32 (7.4%)		
	Grasslands	13 (72.2%)	6 (37.5%)		
	Forests*	83 (36.9%)	31 (28.7%)		
	Grasslands*	13 (72.2%)	5 (100%)		

Wild boars were reported from 36 protected areas (38.7% - Fig. 1), 26 strictly protected (27.9%), and ten areas of sustainable use (27.7%). Wild boars were first cited in protected areas in the studied region in 2005 (two reports), and the most recent invasion was recorded in 2020 (Table 4). The oldest tree detection (two in 2005 and one in 2006) are in the Forests biome, been 2007 the first year with records in the Grasslands biome. Regarding the administrative sphere, in RS, after a rapid expansion, detection remained continuous until 2019, with one or two new records in PAs per year, with a gap of two years (2009 and 2010), with a peak in 2015 with four records. For SC, there is a time gap after 2006 being detected in a new PA only in 2014 until 2017, and there are no new detections until 2020. In the case of PR, the first detections are more spread over the years compared to other states.




Figure 1. Presence of wild boar and control actions. Presence and absence of wild boar and control campaigns in protected areas in southern Brazil.

Table 4. First wild boar records in the PAs. First record of wild boar in environmental protection areas. RS (Rio Grande do Sul), SC (Santa Carina), PR (Paraná). F (Federal), S (State), M (Municipal). Years belonging to the 20th century.

	Year 20th	05	06	07	08	09-10	11	12	13	14	15	16	17	18	19	20
RS	F	1		5		-	1									
	S			1	1	-			1		2		1	1		
	M					-		1			2			1	1	
SC	F	1	1			-				2		1				
	S					-					1		2			
	M					-										
PR	F				1	-					1	1			1	
	S					-					2					1
	M					-										

Wild boars are managed in 13 (36.1%) of the PAs invaded (Fig. 2) – 11 (42.3%) Strictly Protected Areas and two (20%) Sustainable Use Areas; six (37.5%) federals and seven (46.7%) states; or three (50%) in Grasslands and ten (33.3%) in Forests biomes. When asked about the relative importance of wild boar's negative effects against other drivers (in percentage terms), the respondents from the PAs that do not carry out the management of the wild boar was, on average, 28.9% (SD 20.6%), while those that practice the management actions had an average of 59.4% (SD 19.3%). When asked about the relevance of wild boar among other invasive species, the respondents from PAs without management actions had an average score of 40.7% (SD 23.3%), while those that managed wild boar had an average score of 62.6% (SD 21.5%).

The answers from respondents of both PA categories (Strictly Protected and Sustainable Use) were similar (Table 5). Sixteen out of the 22 PAs reported not managing wild boars, justifying it with answers related to low Invasion Intensity. We grouped in this category the answers reported as not a priority, few recorded effects, low number of registered individuals, and recently observed. Nine respondents claimed low management capacity. We grouped in this category the answers reporting lack of technical team or/and structure, lack of formal protocols, bureaucratic difficulties to carry out management, conflicts of interest with the surrounding community, and management plans still in preparation (Fig. 3 and Table 5). "Low number of recorded individuals" and "Lack of technical team/structure" were the most prevalent answers for not carrying out wild boar management. The first reason was indicated by 12 PAs (54.5% of all respondents and 75% of those from the Invasion Intensity group) and was the sole reason reported by nine of them. The second option was informed seven times (31.8% of all respondents and 77.8% of those in the Management Capacity group), and three of them reported the sole reason.




Figure 2. Presence of wild boar in protected areas separated by category of use or Biome. Presence or absence of wild boar in Protected Areas separated by (A) category of use, or (B) by Biome, together with the presence or absence of control actions. In A orange colors refer to Strictly Protected Areas (SP), yellow colors refer to Sustainable Use Areas (SU), and the gray color groups the two categories. In B green colors refer to Tropical and Subtropical Moist Broadleaf Forests (Forests), yellow colors refer to Tropical and Subtropical Grasslands, Savannas and Shrublands (Grasslands) and the gray color groups the two biomes.




Figure 3. Justifications for not carrying out management campaigns. Grouping of protected areas according to subgroups of justifications for not management wild boar. On the right are the subgroups related to Management Capacity and on the left are the subgroups related to Invasion Intensity. The number inside and the thickness of the sphere represent the number of PAs that used the justification. The PAs were only grouped if they used the same justifications.

All the options available in the questionnaire of techniques for wild boar management were selected in the answers (Table 5). Cage trap was the commonest technique used (64.3%) while hunting with dogs was the least reported one (14.3%). The usage pattern is the same when considered between the two types of PAs. On average, 3.5 cages are used in management, showing a difference when separated by type of PA, where Strictly Protected Areas have an average of 2.2 while 5.6 for Sustainable Use Areas. The cage sizes reported vary among the PAs having from two to 32 meters in volume with an average of nine meters. Corrals, varying from one to six, were used in seven PAs. Corn was used as bait in all PAs, whether as coarse grain, on the cob, fermented, or salted (Table 5). Bait is usually used at the entrance and center of the trap and either placed one to seven days before the onset of captures or offered continuously. Cages were placed several weeks before the start of baiting or trapping. In most cases (7 out of 11 PAs), the corral or cages were built by PAs' employees or donated by partner institutions. All PAs that manage wild boar employed irregular efforts without precise seasonal distribution (Table 5).

Regarding external support in the elaboration or execution of management actions, ten PAs were reported to have support from external institutions: four from universities, four from other governmental institutions, and two from NGOs (Table 5). Seven of nine PAs donate the carcasses to hunters if interested, and the other two bury the carcasses at licensed sites. Five PAs did not answer this question. Zoonoses monitoring is performed by only five (out of 14) PAs (Table 3).

Table 5. Summary of questionnaire responses. The table summarizes the answers obtained from the questionnaire. When a PA uses only one action, justification, or method in multiple choice questions, the answer is highlighted in the "Exclu." column. Values with an asterisk are the average of the response values. ¹ Textual responses were grouped into categories according to similarity. SP = Strictly Protected Areas, SU = Sustainable Use Areas.

First section	SP	SU	Total	
- Presence of wild boar				
Yes	27 (27.3%)	12 (32.4%)	39 (27.9%)	
No	72 (72.7%)	25 (67.6%)	97 (72.1%)	
- Wild boar as a generator of adverse effects among all the factors				
With control actions	60.4%*	56.7%*	59.4%*	
Without control actions	31.1%*	25.2%*	28.9%*	
- Wild boar as a generator of adverse effects among invasive species				
With control actions	59.8%*	70%*	62.6%*	
Without control actions	44.6%*	33%*	40.7%*	
Second section	SP	SU	Total	Exclu.
- Performed control actions				
Yes	11 (42.3%)	3 (25%)	14 (36.8%)	
No	15 (57.7%)	9 (75%)	24 (63.2%)	
- Explain the reason why no wild boar control actions were taken ¹				
Invasion Intensity	10 (71.4%)	6 (75%)	16 (72.7%)	11
Not a priority	2 (14.3%)	1 (12.5%)	3 (13.6%)	1
Few recorded effects	1 (7.1%)	2 (25%)	3 (13.6%)	
Low number	7 (50%)	5 (62.5%)	12 (54.5%)	9
Recently observed	1 (7.1%)	0 (0%)	1 (4.5%)	1
Management Capacity	6 (42.9%)	3 (37.5%)	9 (40.9%)	4

Lack of technical team	4 (28.6%)	3 (37.5%)	7 (31.8%)	3
Lack of formal protocols	1 (7.1%)	1 (12.5%)	2 (9.1%)	
Bureaucratic difficulty	2 (14.3%)	0 (0%)	2 (9.1%)	
Conflict of interest	1 (7.1%)	0 (0%)	1 (4.5%)	
Management plan being prepared	2 (14.3%)	0 (0%)	2 (9.1%)	1
- Technical support				
Yes	8 (72.7%)	2 (66.7%)	10 (71.4%)	
No	3 (27.3%)	1 (33.3%)	4 (28.6%)	
- This technical support is provided by:				
University	3	1	4	1
Government agencies	4	0	4	1
NGOs	1	1	2	2
Companies	0	0	0	0
Others	3	1	4	2
- Control technique				
Hunting	5 (45.5.4%)	2 (100%)	7 (53.8%)	
Hunting with dogs	1 (9.1%)	1 (50%)	2 (15.4%)	
Hunting without dogs	4 (36.4%)	2 (100%)	6 (46.2%)	1
Stands	3 (27.3%)	2 (100%)	5 (38.5%)	
Traps	8 (72.7%)	2 (100%)	10 (76.9%)	
Corral	6 (54.6%)	1 (50%)	7 (53.8%)	1
Cage	6 (54.6%)	2 (100%)	8 (61.5%)	1
Fencing	1 (9.1%)	1 (50%)	2 (15.4%)	
Others	2 (18.2%)	0 (0%)	2 (15.4%)	1
- How many cages were used?	2.2*	5.6*	3.5*	
- How many corrals were used?	2.2*	4.5*	2.9*	
- Bait type				
Unspecified corn	4	1	5	3
Fermented corn	1	2	3	1
Dry corn	2	1	3	2
Corn cob	2	-	2	1
Domestic animal carcasses	-	1	1	
Sweet potato	-	1	1	
Leftover vegetables	-	1	1	
Coarse salt	1	1	2	
Third section	SP	SU	Total	Exclu.
- How many individuals were captured?	136.3*	69.2*	91.6*	
- Zoonosis monitoring				
Yes	3 (27.3%)	2 (66.7%)	5 (35.7%)	
No	8 (72.7%)	1 (33.3%)	9 (64.3%)	
- Carcass destination				
Discarded	4	2	6	2
Donated	4	3	7	3
Not Mentioned	5	0	5	

Discussion

In this study, we show that despite the long presence of wild boar in southern Brazil, controlling wild boars in PAs is not a frequent priority, either because its effects are not apparent or because of a lack of resources. We also show that the techniques used follow world standards (trapping and hunting) and corn as bait (Geisser and Reyer 2004; West et al. 2009; VerCauteren et al. 2019; Keuling et al. 2021). However, there is no standard in the effort size or spatial and temporal distribution of managing campaigns.

Despite the wild boar being present in the region since the early 1990s (Hegel et al. 2022), our sample's first reported sightings in PAs were in 2005 for SC and RS and in 2008 for PR. Reinforcing the history of expansion in RS (Hegel et al. 2022), there was a significant advance in the presence between the years 2005 and 2008. First, wild boar detections were in the federal PAs, later in the state and municipal ones, despite the lower number of federal PAs. The lack of detection in municipal areas may be associated with their location, as most are close to urban or peri-urban areas where wild boar tends to avoid areas with higher human presence (Amendolia et al. 2019; Morais et al. 2019) or due to smaller size, by chance are less prone to colonization when compared to federal PAs (Burns 2015; Gallardo et al. 2017). Looking at the use categories, more Sustainable Use PAs reported having wild boar proportionally compared to Strictly Protected Areas. However, more studies are needed to understand if and what may influence this pattern.

Management actions are applied in less than half of the PAs, and it is possible to notice a difference between the categories where more actions are performed in Strictly Protected Areas. This difference may be related to the distribution of resources or the purpose of each type of PA. Nevertheless, when asked about the relevance of wild boar invasion among all factors causing negative effects within the PA, it is not possible to see any difference in the perceived importance of the wild boar effect between the two types of PA. Thus, it is not a matter of perception about the influence of negative effects caused by wild boar. It is only noticeable between those PAs that manage and those that do not manage the species.

Among the reasons for not carrying out the management, the recurrent allegation based on restricted management capacity within individual PAs is compelling evidence for the need for a coordinated control program encompassing the mosaic of federal, state, municipal, and private PAs in a given region, as is being done for managing other biodiversity threats (Miranda et al. 2020; Faria et al. 2022). For successful management, in the case of wild boar, continuous actions are necessary to reduce populations to the point that the effects are within acceptable levels according to the PA conservation objectives (West et al. 2009; VerCauteren et al. 2019; Jori et al. 2021). However, for these campaigns to be efficient, more than control policies are needed, but rather resources to put them into practice. It is not new that the PAs in Brazil are not well-financed, managed, or equipped (Chiaravalloti et al. 2015; de Oliveira and Bernard 2017), and to make matters worse in recent years, environmental policies have been harmed (Bernard et al. 2014; Dobrovolski et al. 2018; Abessa et al. 2019; Golden Kroner et al. 2019), making it even more challenging to face biological invasions. A possible idea would be to share equipment such as corrals and cages between nearby PAs, as animals tend to learn to avoid traps, requiring spatial and temporal rotation. An idea to minimize the losses would be sharing equipment such as corrals and cages between nearby PAs, as animals tend to learn to avoid traps, requiring spatial and temporal rotation (Parkes et al. 2010; Massei et al. 2011; Gaskamp et al. 2021). In addition, exchanging experience between managers,

rangers, and the partner community would increase control efficiency by sharing what went right and wrong in each PA (Meyerson and Mooney 2007; Simpson et al. 2009).

Furthermore, downgrading the National Action Plan (BRASIL 2017) to a state scale could promote the plan's effectiveness as many problems may be at this level as these are at different stages in the wild boar invasion process (Estévez et al. 2015; Early et al. 2016; Courchamp et al. 2017; Shackleton et al. 2019; Hegel et al. 2022). Leaving it in charge of the national sphere to organize and resolve the doubts of managers regarding legislation and protocols since the PAs are divided into two large groups and several subgroups that present different possibilities and restrictions, and the lack of protocols and bureaucratic difficulties were justifications for not carrying out the management in addition to creating means for exchanging experiences between state environmental agencies. The state sphere can oversee organizing the exchange of experience between the agents directly involved (managers, rangers, among others), as they are the ones who plan, organize, and execute the actions within the respective PAs. Thus, it may affect the lack of equipment, people, and conflicts by increasing the efficiency of control campaigns.

An interesting point that deserves an awareness campaign is the justification for "no management" based on the low number of recorded animals in the PA. There is accumulating evidence that the first stages of invasion are the best window for effective control (Allendorf and Lundquist 2003; Mooney et al. 2005; Keiter and Beasley 2017; Giakoumi et al. 2019; Reaser et al. 2020; Ziller et al. 2020), where hunting and trapping would be indicated to keep populations at low levels (Gürtler et al. 2018; VerCauteren et al. 2019; Jori et al. 2021). Most of the PAs within this group had the first record of wild boars very recently (after 2015), contrasting with PAs that manage the species where the first record was prior to 2011. This situation could indicate that populations are still expanding in some areas, which could become a problem in the future (Hegel et al. 2022).

Most PAs that manage wild boar use more than one control technique, which is seen as favorable in management effectiveness assessments (Veitch and Clout 2002; Cruz et al. 2005; Mccann and Garcelon 2008; Parkes et al. 2010; Massei et al. 2011). Cages and corrals are highly effective in controlling wild boar populations (Choquenot et al. 1993), and both techniques are used worldwide to control the species within protected areas and rural environments. The use of corrals has been encouraged in Brazil by some initiatives in the Pampa biome, with construction and use guides (Coelho et al. 2018). Unlike the cage, this technique allows the capture of the whole sounder, which is desirable in population control (Choquenot et al. 1993).

Hunting without dogs was the most used active control technique. Unlike the use of corrals, this technique is intended to remove animals that avoid the traps and should be used as a complement to trapping because they have low effectiveness (West et al. 2009; Hanson et al. 2009). In the Brazilian scenario, the most used technique is hunting with dogs (Rosa et al. 2018), although only two PAs reported using dogs. The use of hunters within PAs to manage wild boar is a topic that needs more debate between environmental managers and the community. In Brazil, the ban on hunting, especially within protected areas, has generated a culture of conflict where those responsible for PAs actively spend resources to prevent hunters from invading the boundaries of areas to kill protected animals or collect endangered plants (Kauano et al. 2017; Ruas et al. 2017; Bragagnolo et al. 2019; Castilho et al. 2019). However, the situation of the wild boar, the only animal

for which hunting is allowed, requires a change in the history of fighting hunting. The PAs do not have the human resources to handle the control activities alone; however, the loss of confidence in hunters due to the Brazilian history makes the partnership between the two difficult.

Corn was the main bait, according to procedures adopted in other countries, since it proved effective (West et al. 2009). However, there are reports that corn may not be attractive depending on the region due to more valuable resources available (West et al. 2009; Hygnstrom et al. 2014). This may happen, for example, in south Brazilian highland forests with abundant *Araucaria angustifolia* seeds, a resource highly consumed by wild boars during winter months (Cervo and Guadagnin 2020). Ideally, bait selection (single species or multispecies) should be locally tested before starting control campaigns (Ballari et al. 2015).

The destination of the carcasses follows the national legislation, exempting the PAs from possible complications (IBAMA 2020). Monitoring diseases could be further encouraged since the wild boar is a species known to be a reservoir of multiple farm animal and human diseases (Maciel et al. 2018; Kmetiuk et al. 2019).

Despite the effort to obtain contact information for all PAs, only 42.3% were accessible, with a return rate of 45.1%. Our sample highly represented federal and state PAs, but less about municipal or private PAs is known. Since this latter comprises the most significant number of PAs, although, in territorial extension, they are smaller, and tend to be managed independently, one way to increase their protection or recovery from wild boar invasions may be recognizing them within multijurisdictional PA network action plans.

Conclusions

In general, we see that most PAs did not detect the species, possibly due to the actual absence of individuals or the lack of human resources to survey the areas. Since the species is still expanding, we believe that some PAs may not have been detected yet due to the presence of a few individuals. We see that management is carried out at the federal and state levels, using similar techniques but with different materials, which can affect the effectiveness of actions. Another point is management campaigns; some carry out continuous management throughout the year, and others punctually. This is a possible reflection of the lack of an adaptive management plan with well-defined objectives and goals. This does not detract from the merits of the initiatives. However, it is not effective in terms of population control. Thus, an organization of environmental agencies is necessary to assist environmental managers in creating well-defined plans that go in the direction of the national plan.

Bibliography

- Abessa D, Famá A, Buruaem L (2019) The systematic dismantling of Brazilian environmental laws risks losses on all fronts. *Nat Ecol Evol* 3:510–511. <https://doi.org/10.1038/s41559-019-0855-9>
- Allendorf FW, Lundquist LL (2003) Introduction: Population Biology, Evolution, and Control of Invasive Species. *Conservation Biology* 17:24–30

Amendolia S, Lombardini M, Pierucci P, Meriggi A (2019) Seasonal spatial ecology of the wild boar in a peri-urban area. *Mamm Res* 64:387–396. <https://doi.org/10.1007/s13364-019-00422-9>

Andreu J, Vilà M, Hulme PE (2009) An Assessment of Stakeholder Perceptions and Management of Noxious Alien Plants in Spain. *Environmental Management* 43:1244–1255. <https://doi.org/10.1007/s00267-009-9280-1>

Ballari SA, Cuevas MF, Ojeda RA, Navarro JL (2015) Diet of wild boar (*Sus scrofa*) in a protected area of Argentina: the importance of baiting. *Mamm Res* 60:81–87. <https://doi.org/10.1007/s13364-014-0202-0>

Barrios-Garcia MN, Ballari SA (2012) Impact of wild boar (*Sus scrofa*) in its introduced and native range: a review. *Biol Invasions* 14:2283–2300. <https://doi.org/10.1007/s10530-012-0229-6>

Bernard E, Penna L a. O, Araújo E (2014) Downgrading, Downsizing, Degazettement, and Reclassification of Protected Areas in Brazil. *Conservation Biology* 28:939–950. <https://doi.org/10.1111/cobi.12298>

Bragagnolo C, Gama GM, Vieira FAS, et al (2019) Hunting in Brazil: What are the options? *Perspectives in Ecology and Conservation* 17:71–79. <https://doi.org/10.1016/j.pecon.2019.03.001>

BRASIL PJ (2017) Plano Nacional de Prevenção, Controle e Monitoramento do Javali (*Sus scrofa*) no Brasil.

Burns KC (2015) A Theory of Island Biogeography for Exotic Species. *The American Naturalist* 186:441–451. <https://doi.org/10.1086/682934>

Butchart SHM, Walpole M, Collen B, et al (2010) Global Biodiversity: Indicators of Recent Declines. *Science* 328:1164–1168. <https://doi.org/10.1126/science.1187512>

Castilho LC, De Vleeschouwer KM, Milner-Gulland EJ, Schiavetti A (2018) Attitudes and Behaviors of Rural Residents Toward Different Motivations for Hunting and Deforestation in Protected Areas of the Northeastern Atlantic Forest, Brazil. *Tropical Conservation Science* 11:1940082917753507. <https://doi.org/10.1177/1940082917753507>

Castilho LC, Vleeschouwer KMD, Milner-Gulland EJ, Schiavetti A (2019) Hunting of mammal species in protected areas of the southern Bahian Atlantic Forest, Brazil. *Oryx* 53:687–697. <https://doi.org/10.1017/S0030605317001247>

Cervo IB, Guadagnin DL (2020) Wild boar diet and its implications on agriculture and biodiversity in Brazilian forest–grassland ecoregions. *Anim Biodiv Conserv* 123–136. <https://doi.org/10.32800/abc.2020.43.0123>

Chiaravalloti RM, Delelis C, Tofoli C, et al (2015) Federal protected areas management strategies in Brazil: sustainable financing, staffing, and local development. *Natureza & Conservação* 13:30–34. <https://doi.org/10.1016/j.ncon.2015.05.003>

Choquenot D, Kilgour RJ, Lukins BS (1993) An evaluation of feral pig trapping. *Wildl Res* 20:15–21. <https://doi.org/10.1071/wr9930015>

Coelho R, Jacques J, Molino N, et al (2018) Controle de porcos ferais – Javalis, Construção de jaula curral modelo Pampa, Guia para o Produtor Rural. Instituto Chico Mendes para a Conservação da Biodiversidade, Ministério do Meio Ambiente.

Courchamp F, Fournier A, Bellard C, et al (2017) Invasion Biology: Specific Problems and Possible Solutions. *Trends in Ecology & Evolution* 32:13–22. <https://doi.org/10.1016/j.tree.2016.11.001>

Cruz F, Josh Donlan C, Campbell K, Carrion V (2005) Conservation action in the Galàpagos: feral pig (*Sus scrofa*) eradication from Santiago Island. *Biological Conservation* 121:473–478. <https://doi.org/10.1016/j.biocon.2004.05.018>

de Oliveira APC, Bernard E (2017) The financial needs vs. the realities of in situ conservation: an analysis of federal funding for protected areas in Brazil's Caatinga. *Biotropica* 49:745–752. <https://doi.org/10.1111/btp.12456>

de Souza JB, Alves RRN (2014) Hunting and Wildlife use in an Atlantic Forest Remnant of Northeastern Brazil. *Tropical Conservation Science* 7:145–160. <https://doi.org/10.1177/194008291400700105>

Deberdt AJ, Scherer SB (2007) The wild boar: occurrence and management of the species in Brazil. *Natureza & Conservação* 5:101–114

Dinerstein E, Olson D, Joshi A, et al (2017) An Ecoregion-Based Approach to Protecting Half the Terrestrial Realm. *BioScience* 67:534–545. <https://doi.org/10.1093/biosci/bix014>

Dobrovolski R, Loyola R, Rattis L, et al (2018) Science and democracy must orientate Brazil's path to sustainability. *Perspectives in Ecology and Conservation* 16:121–124. <https://doi.org/10.1016/j.pecon.2018.06.005>

Dudley N, Stolton S (2010) *Arguments for Protected Areas: Multiple Benefits for Conservation and Use*. Routledge

Early R, Bradley BA, Dukes JS, et al (2016) Global threats from invasive alien species in the twenty-first century and national response capacities. *Nat Commun* 7:12485. <https://doi.org/10.1038/ncomms12485>

Estévez RA, Anderson CB, Pizarro JC, Burgman MA (2015) Clarifying values, risk perceptions, and attitudes to resolve or avoid social conflicts in invasive species management. *Conservation Biology* 29:19–30. <https://doi.org/10.1111/cobi.12359>

Faria L, Carvalho BM de, Carneiro L, et al (2022) Invasive species policy in Brazil: a review and critical analysis. *Environmental Conservation* 1–6. <https://doi.org/10.1017/S0376892922000406>

Foxcroft LC, Pyšek P, Richardson DM, et al (2017) Erratum to: Plant invasion science in protected areas: progress and priorities. *Biol Invasions* 19:2503–2505. <https://doi.org/10.1007/s10530-017-1445-x>

Fryxell JM, Sinclair ARE, Caughley G (2014) *Wildlife Ecology, Conservation, and Management*. John Wiley & Sons

Gallardo B, Aldridge DC, González-Moreno P, et al (2017) Protected areas offer refuge from invasive species spreading under climate change. *Global Change Biology* 23:5331–5343. <https://doi.org/10.1111/gcb.13798>

Gaskamp JA, Gee KL, Campbell TA, et al (2021) Effectiveness and Efficiency of Corral Traps, Drop Nets and Suspended Traps for Capturing Wild Pigs (*Sus scrofa*). *Animals* 11:1565. <https://doi.org/10.3390/ani11061565>

Geisser H, Reyer H-U (2004) Efficacy of Hunting, Feeding, and Fencing to Reduce Crop Damage by Wild Boars. *The Journal of Wildlife Management* 68:939–946. [https://doi.org/10.2193/0022-541X\(2004\)068\[0939:EOHFAF\]2.0.CO;2](https://doi.org/10.2193/0022-541X(2004)068[0939:EOHFAF]2.0.CO;2)

Genovesi P, Monaco A (2013) Guidelines for Addressing Invasive Species in Protected Areas. In: Foxcroft LC, Pyšek P, Richardson DM, Genovesi P (eds) *Plant Invasions in Protected Areas: Patterns, Problems and Challenges*. Springer Netherlands, Dordrecht, pp 487–506

Georgakopoulou A (2019) *Discourse Analysis: An Introduction*. Edinburgh University Press

Giakoumi S, Katsanevakis S, Albano PG, et al (2019) Management priorities for marine invasive species. *Science of The Total Environment* 688:976–982. <https://doi.org/10.1016/j.scitotenv.2019.06.282>

Golden Kroner RE, Qin S, Cook CN, et al (2019) The uncertain future of protected lands and waters. *Science* 364:881–886. <https://doi.org/10.1126/science.aau5525>

Gürtler RE, Rodríguez-Planes LI, Gil G, et al (2018) Differential long-term impacts of a management control program of axis deer and wild boar in a protected area of north-eastern Argentina. *Biol Invasions* 20:1431–1447. <https://doi.org/10.1007/s10530-017-1635-6>

Hanson LB, Mitchell MS, Grand JB, et al (2009) Effect of experimental manipulation on survival and recruitment of feral pigs. *Wildl Res* 36:185–191. <https://doi.org/10.1071/WR08077>

Hegel CGZ, Faria GMM, Ribeiro B, et al (2022) Invasion and spatial distribution of wild pigs (*Sus scrofa* L.) in Brazil. *Biol Invasions*. <https://doi.org/10.1007/s10530-022-02872-w>

Hygnstrom SE, Timm RM, Curtis PD, et al (2014) *Prevention and Control of Wildlife Damage*, 5th Edition. *Proceedings of the Vertebrate Pest Conference* 26:. <https://doi.org/10.5070/V426110401>

IBAMA (2020) *Manual de boas práticas para o controle de javali*. Grazielle Oliveira Batista, Brasília, DF.

Jaureguiberry P, Titeux N, Wiemers M, et al (2022) The direct drivers of recent global anthropogenic biodiversity loss. *Science Advances* 8:eabm9982. <https://doi.org/10.1126/sciadv.abm9982>

Jori F, Massei G, Licoppe A, et al (2021) 8. Management of wild boar populations in the European Union before and during the ASF crisis. In: *Understanding and combatting African Swine Fever*. Wageningen Academic Publishers, pp 197–228

Kauano ÉE, Silva JMC, Michalski F (2017) Illegal use of natural resources in federal protected areas of the Brazilian Amazon. *PeerJ* 5:e3902. <https://doi.org/10.7717/peerj.3902>

Keiter DA, Beasley JC (2017) Hog Heaven? Challenges of Managing Introduced Wild Pigs in Natural Areas. *naar* 37:6–16. <https://doi.org/10.3375/043.037.0117>

Keuling O, Strauß E, Siebert U (2016) Regulating wild boar populations is "somebody else's problem"! - Human dimension in wild boar management. *Science of The Total Environment* 554–555:311–319. <https://doi.org/10.1016/j.scitotenv.2016.02.159>

Keuling O, Strauß E, Siebert U (2021) How Do Hunters Hunt Wild Boar? Survey on Wild Boar Hunting Methods in the Federal State of Lower Saxony. *Animals* 11:2658. <https://doi.org/10.3390/ani11092658>

Kiringe JW, Okello MM, Ekajul SW (2007) Managers' perceptions of threats to the protected areas of Kenya: prioritization for effective management. *Oryx* 41:314–321. <https://doi.org/10.1017/S0030605307000218>

Klinger T, Padilla DK, Britton-Simmons K (2006) Two invaders achieve higher densities in reserves. *Aquatic Conservation: Marine and Freshwater Ecosystems* 16:301

Kmetiuk LB, Krawczak FS, Machado FP, et al (2019) Ticks and serosurvey of anti-Rickettsia spp. antibodies in wild boars (*Sus scrofa*), hunting dogs and hunters of Brazil. *PLOS Neglected Tropical Diseases* 13:e0007405. <https://doi.org/10.1371/journal.pntd.0007405>

La Sala LF, Burgos JM, Caruso NC, et al (2023) Wild pigs and their widespread threat to biodiversity conservation in South America. *Journal for Nature Conservation* 73:126393. <https://doi.org/10.1016/j.jnc.2023.126393>

Laurance WF, Carolina Useche D, Rendeiro J, et al (2012) Averting biodiversity collapse in tropical forest protected areas. *Nature* 489:290–294. <https://doi.org/10.1038/nature11318>

Liu X, Blackburn TM, Song T, et al (2020) Animal invaders threaten protected areas worldwide. *Nat Commun* 11:2892. <https://doi.org/10.1038/s41467-020-16719-2>

Long JL (2005) Introduced Mammals of the World: Their History, Distribution and Influence. John Long. A Review by Robert L Close. *Aust Mammalogy* 27:109–110. https://doi.org/10.1071/am05109_br

Maciel ALG, Loiko MR, Bueno TS, et al (2018) Tuberculosis in Southern Brazilian wild boars (*Sus scrofa*): First epidemiological findings. *Transboundary and Emerging Diseases* 65:518–526. <https://doi.org/10.1111/tbed.12734>

Massei G, Roy S, Bunting R (2011) Too many hogs?: A review of methods to mitigate impact by wild boar and feral hogs. *Human-Wildlife Interactions* 5:79–99

Mccann BE, Garcelon DK (2008) Eradication of Feral Pigs From Pinnacles National Monument. *The Journal of Wildlife Management* 72:1287–1295. <https://doi.org/10.2193/2007-164>

Meyerson LA, Mooney HA (2007) Invasive alien species in an era of globalization. *Frontiers in Ecology and the Environment* 5:199–208. [https://doi.org/10.1890/1540-9295\(2007\)5\[199:IASIAE\]2.0.CO;2](https://doi.org/10.1890/1540-9295(2007)5[199:IASIAE]2.0.CO;2)

Miranda RJ, Nunes JACC, Creed JC, et al (2020) Brazil policy invites marine invasive species. *Science* 368:481–481. <https://doi.org/10.1126/science.abb7255>

- Mooney HA, Mack R, McNeely JA, et al (2005) *Invasive Alien Species: A New Synthesis*. Island Press
- Morais TA, Rosa CA, Azevedo CS, et al (2019) Factors affecting space use by wild boars (*Sus scrofa*) in high-elevation tropical forests. *Can J Zool* 97:971–978. <https://doi.org/10.1139/cjz-2019-0130>
- Parkes JP, Ramsey DSL, Macdonald N, et al (2010) Rapid eradication of feral pigs (*Sus scrofa*) from Santa Cruz Island, California. *Biological Conservation* 143:634–641. <https://doi.org/10.1016/j.biocon.2009.11.028>
- Pyšek P, Hulme PE, Simberloff D, et al (2020) Scientists' warning on invasive alien species. *Biological Reviews* 95:1511–1534. <https://doi.org/10.1111/brv.12627>
- Reaser JK, Burgiel SW, Kirkey J, et al (2020) The early detection of and rapid response (EDRR) to invasive species: a conceptual framework and federal capacities assessment. *Biol Invasions* 22:1–19. <https://doi.org/10.1007/s10530-019-02156-w>
- Ren J, Chen J, Xu C, et al (2021) An invasive species erodes the performance of coastal wetland protected areas. *Science Advances* 7:eabi8943. <https://doi.org/10.1126/sciadv.abi8943>
- Risch DR, Ringma J, Price MR (2021) The global impact of wild pigs (*Sus scrofa*) on terrestrial biodiversity. *Sci Rep* 11:13256. <https://doi.org/10.1038/s41598-021-92691-1>
- Rosa CAD, Wallau MO, Pedrosa F (2018) Hunting as the main technique used to control wild pigs in Brazil. *Wildlife Society Bulletin* 42:111–118. <https://doi.org/10.1002/wsb.851>
- Ruas RMS, Furtado DC, Guerra GAD, et al (2017) CAÇA, CAPTURA E USO DA FAUNA SILVESTRE NO BRASIL COMO CRIMES AMBIENTAIS E TABU CIENTÍFICO: REFLEXÃO SOBRE CATEGORIAS TEÓRICAS. *HOLOS* 5:37–54. <https://doi.org/10.15628/holos.2017.5660>
- Rylands AB, Brandon K (2005) Brazilian Protected Areas. *Conservation Biology* 19:612–618. <https://doi.org/10.1111/j.1523-1739.2005.00711.x>
- Sales LP, Ribeiro BR, Hayward MW, et al (2017) Niche conservatism and the invasive potential of the wild boar. *Journal of Animal Ecology* 86:1214–1223. <https://doi.org/10.1111/1365-2656.12721>
- Salvador C (2012) Ecologia e manejo de javali (*Sus scrofa* L.) na América do Sul [Ecology and management of Eurasian wild boar (*Sus scrofa* L.) in South America]
- Sampaio AB, Schmidt IB (2013) Espécies exóticas invasoras em unidades de conservação federais do Brasil. *Biodiversidade Brasileira-BioBrasil* 32–49
- Shackleton RT, Adriaens T, Brundu G, et al (2019) Stakeholder engagement in the study and management of invasive alien species. *Journal of Environmental Management* 229:88–101. <https://doi.org/10.1016/j.jenvman.2018.04.044>
- Simberloff D, Martin J-L, Genovesi P, et al (2013) Impacts of biological invasions: what's what and the way forward. *Trends in Ecology & Evolution* 28:58–66. <https://doi.org/10.1016/j.tree.2012.07.013>
- Simpson A, Jarnevich C, Madsen J, et al (2009) Invasive species information networks: collaboration at multiple scales for prevention, early detection, and rapid response to

invasive alien species. Biodiversity 10:5–13.
<https://doi.org/10.1080/14888386.2009.9712839>

Spear D, Foxcroft LC, Bezuidenhout H, McGeoch MA (2013) Human population density explains alien species richness in protected areas. *Biological Conservation* 159:137–147.
<https://doi.org/10.1016/j.biocon.2012.11.022>

Taylor S (2013) *What is Discourse Analysis?* Bloomsbury Academic

Tu M, Robison RA (2013) Overcoming Barriers to the Prevention and Management of Alien Plant Invasions in Protected Areas: A Practical Approach. In: Foxcroft LC, Pyšek P, Richardson DM, Genovesi P (eds) *Plant Invasions in Protected Areas: Patterns, Problems and Challenges*. Springer Netherlands, Dordrecht, pp 529–547

Veitch CR, Clout MN (2002) *Turning the Tide: The Eradication of Invasive Species : Proceedings of the International Conference on Eradication of Island Invasives*. IUCN

VerCauteren KC, Beasley JC, Ditchkoff SS, et al (2019) *Invasive Wild Pigs in North America: Ecology, Impacts, and Management*. CRC Press

West B, Cooper A, Armstrong J (2009) *Managing Wild Pigs: A Technical Guide*. Human–Wildlife Interactions Monographs. <https://doi.org/10.26079/600a-ed8d>

Ziller SR, Dechoum M de S, Silveira RAD, et al (2020) A priority-setting scheme for the management of invasive non-native species in protected areas. *NeoBiota* 62:591–606.
<https://doi.org/10.3897/neobiota.62.52633>