

PREPRINT

Author-formatted, not peer-reviewed document posted on 01/03/2024

DOI: <https://doi.org/10.3897/arphapreprints.e121861>

Redefining the megagenus *Erica* L. (Ericaceae): the contributions of E. G. H. Oliver and I. M. Oliver (née Nitzsche) to taxonomy and nomenclature

 E. Charles Nelson,  Michael Pirie, Dirk Bellstedt

Disclaimer on biological nomenclature and use of preprints

The preprints are preliminary versions of works accessible electronically in advance of publication of the final version. They are not issued for purposes of botanical, mycological or zoological nomenclature and **are not effectively/validly published in the meaning of the Codes**. Therefore, nomenclatural novelties (new names) or other nomenclatural acts (designations of type, choices of priority between names, choices between orthographic variants, or choices of gender of names) **should NOT be posted in preprints**. The following provisions in the Codes of Nomenclature define their status:

International Code of Nomenclature for algae, fungi, and plants (ICNafp)

Article 30.2: "An electronic publication is not effectively published if there is evidence within or associated with the publication that its content is merely preliminary and was, or is to be, replaced by content that the publisher considers final, in which case only the version with that final content is effectively published." In order to be validly published, a nomenclatural novelty must be effectively published (Art. 32.1(a)); in order to take effect, other nomenclatural acts must be effectively published (Art. 7.10, 11.5, 53.5, 61.3, and 62.3).

International Code of Zoological Nomenclature (ICZN)

Article: 21.8.3: "Some works are accessible online in preliminary versions before the publication date of the final version. Such advance electronic access does not advance the date of publication of a work, as preliminary versions are not published (Article 9.9)".

Redefining the megagenus *Erica* L. (Ericaceae): the contributions of E. G. H. Oliver and I. M. Oliver (née Nitzsche) to taxonomy and nomenclature

Running title: Redefining the megagenus *Erica*

E. CHARLES NELSON ¹, MICHAEL D. PIRIE ² and D. U. BELLSTEDT ³

1 *Tippitiwitchet Cottage, 255A Broadgate, Sutton St Edmund PE12 0LT, UK*
ORCID 0000-0002-3909-4388

2 *University Museum, University of Bergen, Postboks 7800, NO-5020 Bergen, Norway*
ORCID 0000-0003-0403-4470

3 *Dept of Biochemistry, University of Stellenbosch, Stellenbosch, South Africa*
ORCID 0000-0002-6376-4855

Corresponding author: Michael D. Pirie (Michael.Pirie@uib.no)

Abstract

The megagenus *Erica* L. (Ericaceae), as it is recognised today, includes 851 species of evergreen shrubs or small trees, the majority of which are endemic to the Cape Floristic Region of South Africa. From the first descriptions in Linnaeus's *Genera plantarum*, a succession of authors ascribed the steadily accumulating numbers of known species to various of a total of 72 different genera. Until the latter half of the twentieth century, so called 'minor genera' such as *Philippia* Klotzsch and *Blaeria* L. were still recognised for many African species. The now uncontroversial inclusive circumscription of *Erica*, and a substantial proportion of its currently recognised species diversity, was conceptualised, described, and illustrated by the South African botanists E. G. H. ('Ted') Oliver and Inge M. Oliver in a succession of works published from 1964 to the present day. We review the historical development of generic delimitation in *Erica sens. lat.*, focusing on the contribution of the Olivers to the current state of systematic knowledge of the genus, and presenting an overview and complete lists of literature and of taxa that they authored.

Key words: Ericoideae; nomenclature; taxonomy

Introduction: historical conspectus

Erica L. (subfamily Ericoideae, family Ericaceae), as understood today, is one of the largest genera in the Angiospermae. World Flora Online (WFO; www.wfo.org) currently recognizes 852 included species (<https://wfo.plantlist.org/taxon/wfo-4000013772-2023-12>; revised to 851 in Oliver et al. 2024; Elliot et al. in prep.). Within this concept of *Erica*, 72 genera (WFO: 69; plus *Apogandrum* Neck. [wfo-4000043784], *Kolbia* Adans. [wfo-4100003440], and *Lopherina* Juss. [wfo-4000043791]) that were designated and named at various times since 1753 have been subsumed – some of these were known informally as ‘minor genera’ following Phillips’s use of that phrase (Phillips 1944), until the major revision by E. G. H. Oliver ‘sank’ all of them into *Erica* (Oliver 2000a; Oliver & Oliver 2000a).


The taxonomic and nomenclatural history of *Erica* is complicated. Carl Linnaeus’s *Genera plantarum* (Linnaeus 1737: 110) and *Species plantarum* (Linnaeus 1753: 112, 352–356) are the starting points for modern classification and nomenclature (Oliver 2007; Jarvis 2007). In the first edition of *Genera plantarum*, Linnaeus (1737: 110 genus no. 312) defined *Erica* in just 51 Latin words, exclusively referring to the anatomy and morphology of the flowers, placing the genus in Class VIII ‘Octandria Monogynia’ based on the numbers of male (eight stamens) and female (one gynoeceum) organs in each flower. (By way of contrast, the most recent definition of *Erica* (Oliver 2000a) exceeds 350 words (in English) (Appendix 1) (for a more succinct description, about 100 words, see Oliver 2004).)

In the first edition of *Species plantarum* Linnaeus described and named only 23 species of *Erica*. His artificial method of classification led him to nest the widespread, northern-hemisphere shrubby *Calluna vulgaris* L. within *Erica* (as *Erica vulgaris* L.: Linnaeus 1753: 352 no. 1). On the same artificial basis, another *Erica*-like African shrub with only four stamens and a solitary gynoeceum (‘Tetrandria Monogynia’), was placed by him in a separate genus named *Blaeria* L. (as *B. ericoides* L.: Linnaeus 1753: 112). Forty-eight of the binomial names published by Linnaeus in eight works produced during his lifetime – in *Species plantarum* (Linnaeus 1753) and its second edition (Linnaeus 1762), a dissertation defended by Jacob Printz entitled *Plantae rariores Africanae* (Linnaeus 1760), the tenth and twelfth editions of *Systema naturae* (Linnaeus 1759, 1767a), two editions of *Mantissa plantarum* (Linnaeus 1767b, 1771) and in a second dissertation *De Erica* (Linnaeus 1770) defended by Johan Adolph Dahlgren – are still in use in *Erica* (see entries in Jarvis 2007: 497–501). Since Linnaeus’s time, the number of taxa recognized as belonging to *Erica* has multiplied vastly, presenting today’s taxonomists with not a few difficulties in delimiting taxa at generic and subgeneric levels.

Vegetatively, *Erica* species (commonly called heaths or heathers) are relatively similar being shrubs or less frequently small trees with small, evergreen, linear-oblong (‘needle-like’) leaves arranged in whorls. The leaves often have revolute margins that can touch on the underside – this general type of leaf morphology is termed ‘ericoid’. As the individual taxa are so similar in foliage, taxonomists, since Linnaeus’s time, have traditionally relied on floral characteristics to demarcate subgeneric taxa and distinguish between species, rather than foliage morphology (for example, see Linnaeus 1770) (Fig. 1). Linnaeus (1753: 352, 354) used the presence/absence of a pair of awns at the base of each anther to subdivide the species he knew – ten species with awned anthers (‘*Antheris bicornibus*’) were separated from thirteen species with muticous (without awns) anthers (‘*Antheris [sic] simplicibus obtusis*’).

emarginatis'). With additional species to accommodate, this scheme was modified to separate those species with included stamens – stamens that did not protrude beyond the mouth of the urn-shaped or tubular corolla – from others with exserted stamens (Linnaeus 1770) (Fig. 1). The number and morphology of the stamens in the flowers of shrubs that could be recognized as *Erica*-like continued to dominate *Erica* taxonomy until the end of the twentieth century. Additional characters linked with the morphology of the gynoecium, particularly the number of locules (ranging from one to eight) comprising the ovary and the number of ovules per locule, and whether the mature capsule was indehiscent or dehiscent, were employed in generic definitions.

Fig. 1. Flowers of *Erica* species known to Carl Linnaeus, from Linnaeus (1770).




The eighteenth-century and nineteenth-century botanists who worked to identify, name, and describe *Erica sens. lat.* species and who attempted to subdivide the numerous species into discrete genera were almost all based in Europe. A few had travelled and collected in Africa. Thus, most of their work entailed examining a relatively small number of pressed and dried specimens. Access to living material was largely limited to the species indigenous in the northern hemisphere, and to the relatively small number of cultivated plants maintained in European gardens. While there was a ‘mania’ for cultivating southern African *Erica* (the so-called ‘Cape heaths’) particularly during the first part of the nineteenth century (Nelson and Oliver 2004; Nelson and Pirie 2022), the plants grown were neither representative of the genus throughout its geographical range nor of its complex morphology. Small-flowered wind-pollinated heathers, for example, were not fashionable and did not attract gardeners. Besides, artificial and accidental cross-pollination in cultivation had produced a plethora of hybrids that tended to be preferred by horticulturists. The major attempts to produce monographs about *Erica* were thus both incomplete and compromised.

The only universal treatment available before 1900 was George Bentham’s contribution to the seventh volume of Agustin Pyramus de Candolle’s *Prodromus systematis naturalis regni vegetabilis* published in 1839 (Bentham 1839; see also Nelson 2005). In late 1906, when the Ericaceae section (parts 1–3) of the fourth volume of *Flora Capensis* (Bolos et al. 1905–1906), covering the plants of southern Africa, was completed, an integrated treatment of diverse *Erica* flora of the Cape Floristic Region became available. This treatment of *Erica* (in a restricted sense) had been produced by Francis Guthrie (1831–1899) and Harry Bolus (1834–1911) (with assistance latterly from his future daughter-in-law, Harriet Margaret Louisa Kensit (*olim* Bolus) (1877–1970)), botanists who lived in the Cape Province of South Africa where *Erica* species are indigenous and numerous. Regarding their concept of *Erica*, Guthrie & Bolus (1906: 5) commented that ‘The genus is remarkable for an unusual degree of variability in the form of almost all its organs. It is therefore one difficult of definition as to its species and of arrangement into satisfactory natural groups.’ They had subdivided the 469 southern African species of *Erica* that they recognized into five subgenera and 41 sections, as well as accepting that close to 160 other species should be placed outside *Erica* in 22 ‘minor genera’ (Phillips 1944: 69; Oliver 2000b: 55). These ‘minor genera’, six being monotypic, were treated by Nicholas Edward Brown (1849–1934), a botanist based at the Royal Botanic Gardens, Kew (Brown 1906). However, many other species native elsewhere in tropical and subtropical Africa and on Atlantic and Indian Ocean islands were not integral in this treatment. Edwin Percy Phillips (1884–1967) was the next to tackle the complexities of the African *Erica sens. lat.* His treatments of Ericaceae (Phillips 1926, 1944, 1951) also include *Vaccinium* L. (Vaccinioideae), represented by a single species now treated as *V. exul* Bolus, which occurs on the Eastern Escarpment (Mpumalanga and Limpopo Provinces), in Swaziland and in Malawi, in habitats above 1,200m altitude (Bester 2015). All other Ericaceae in South Africa represent *Erica sens. lat.* In his paper ‘Notes on the minor genera of Ericaceae’, Phillips (1944) reduced the 22 ‘minor genera’ retained in *Flora Capensis* to six, a scheme he maintained in the second edition of his *The genera of South African flowering plants* (Phillips 1951). Emphasising that he considered that the number of ovules per locule was ‘a more important character than the number of ovary-chambers [locules]’ in any attempt at a natural classification of the South African Ericoideae, Phillips (1944, 1951) retained only *Erica* L., *Blaeria* L., *Eremia* D. Don, *Sympezia* Licht., *Scyphogyne* Brongn., *Salaxis* Salisb. and *Lagenocarpus* Klotzsch.

Taxonomic studies of E. G. H. Oliver and I. M. Oliver

Edward ('Ted') George Hudson Oliver became fascinated by the diversity of Cape flora and especially fynbos vegetation after he enrolled as an undergraduate, to study zoology, at the University of Cape Town in the late 1950s. His attention was soon diverted from animals, and he became 'obsessed' by the Cape heaths because 'they were small and delicate with a seemingly infinite variety of shapes and colours' (Oliver 2000b). He was already collecting *Erica* in the field and making discoveries and, according to Colonel Hugh Arthur Baker (1896–1976) when he named *E. oliveri* (Fig. 2) in July 1962, 'Mr. E. G. H. Oliver ... seems destined to add many more to the 600 or so [*Erica* species] already described' (Baker 1962: 198).

Fig. 2. *Erica oliveri* H.A. Baker, drawn by E. G. H. Oliver (Baker 1962): 'The species has been named after Mr. E. G. H. Oliver who is making a study of the genus and of the minor genera and has already collected a number of hitherto unknown species and seems destined to add many more to the 600 or so [*sic*] already described.'



Following graduation, he undertook a study of *Acrostemon*, one of the ‘minor’ genera, presenting his Master’s thesis in October 1964. In this (Oliver 1964) he noted ‘complete overlap between *Acrostemon* Klotzsch and *Hexastemon* Klotzsch allowing no character for the separation of the two genera’ and therefore proposed to incorporate *Hexastemon* in *Acrostemon*. Likewise, comparing the monotypic genus *Arachnocalyx* Klotzsch, which had been designated by Robert Harold Compton (1886–1979) (Compton 1935), with the single species *Acrostemon viscidus* N.E.Br., Oliver found ‘much closer similarity’ between *Arachnocalyx eriocephalus* (Klotzsch) N.E.Br. and *Acrostemon viscidus* ‘than had previously been suspected’ (Oliver 1964: 105). To gain wider insights into the patterns of morphological similarities and differences, the species then assigned to *Syndesmanthus* Klotzsch and *Simochilus* Klotzsch were investigated too, and again Oliver (1964: 105) noted that ‘a certain degree of intergrading occurs in the differentiating characters between the three genera to an extent where it becomes difficult to distinguish and place species into any one of the genera.’ Discussing the six ‘minor genera’ that Phillips had retained, Oliver (1964: 115) concluded that Phillips’s scheme was ‘completely artificial’ and did not ‘reflect the correct relationships between the genera which [had] been reduced to synonymy. ... With the addition of new and well collected material, a reconsideration of the generic boundaries will certainly be necessary in a future revision ... [T]he problem of generic distinction will have to be worked out carefully and thoroughly.’ At that time, some of the ‘minor genera’ were still regarded as monospecific, including the European endemic *Bruckenthalia* Rchb. and the last African ‘minor’ genus to be described, *Stokoeanthus* E.G.H. Oliv. (Oliver 1976). Commenting on his decision to describe and name that new monotypic genus, he stated (Oliver (1976):

The relationship of [*Stokoeanthus*] appears to me to be with *Erica* and *Blaeria* and to some extent with *Eremia*, but it does not fit into any of them as presently constituted. From *Erica* it differs in having 4 stamens and a 2-celled ovary, from *Blaeria* in having 2 cells to the ovary and from *Eremia* in having 4 stamens and more than 1 ovule per cell. To change the generic limits of any of these genera to force the inclusion of the new taxon would, in my opinion, be impracticable and would cause repercussions in the relationships of and differences between many other genera of the Ericoideae.

Thus, Ted laid down the basis for the work that consumed his time for the next 35 years, culminating in the elimination of all the ‘minor genera’ (Oliver 1987; 1988; 1992; 1993b; 1993a; 1993c; 1994; 1996; 2000a) and the subsuming of all their species into the megagenus *Erica* (Schumann et al. 1992: 244; Oliver 2000a; Oliver & Oliver 2000a). By examining many more plants than had been available to preceding botanists, he noticed clearly overlapping characters in the ‘minor genera’ and apparently discontinuous variation coalescing through various intermediate states, a good example being the capacity of the mature capsule to dehisce.

From about 1974, Oliver was assisted in his work on *Erica*, especially the ‘minor genera’, by Inge Magdalene Nitzsche (1947–2003), who had studied botany and zoology at the University of Cape Town (1967–1969) and also had a diploma in fine arts (1971–1972). They married in February 1974. Inge’s remarkable pen-and-ink drawings of anatomical and morphological details of the species (often not signed) were to be an integral component of papers about *Erica* (and the ‘minor genera’) published from 1976 onwards: early (unsigned) examples of Inge Oliver’s extraordinary illustrations were published in the paper ‘revising’ *Eremia* and *Eremiella* (Oliver 1976: fig. 2, p. 34 *Eremia totta* (Thunb.) D.Don; fig. 9, p. 40

E. curvistyla (N.E.Br.) E.G.H. Oliv.; fig. 14, p. 44 *E. brevifolia* Benth.) (for the eponymous *Erica ingeana* E.G.H. Oliver, see Oliver & Oliver 1991: 140–142 (Fig. 3)). In 1988, a grant enabled Inge to be employed as research assistant in the Bolus Herbarium at the University of Stellenbosch ‘to help with all the technical work – the numerous dissections, drawings and recording of all the details of variations in the plants. This ... helped tremendously with the research and ... Inge’s work also helped with the finalisation of the analyses of the minor ericaceous genera ...’ (Oliver 2000b; Nelson 2004). Ted Oliver submitted his doctoral thesis to the University of Cape Town in 1999 (Oliver 1999) and published the monograph on the ‘minor genera’, integrating all of them into the redefined megagenus *Erica* in 2000. *Field guide to the Ericas of the Cape Peninsula* (Oliver & Oliver 2000b), a handy, pocketable manual, represents another aspect of these collaborative studies, making available an identification aid, illustrated with simple line drawings (Fig. 4), for naturalists in general.

Fig. 3. *Erica ingeana* E.G.H. Oliver, drawn by Inge M. Oliver (Oliver 1991).





FIGURE 4. —*Erica ingeana*. A. flowering branch. $\times 1$. B. branch. C. leaf. D. flower. E. bract. F. sepal. G. anther. front, side and back views. H. gynoecium. B–H. $\times 25$. All drawn from the type, *Oliver 5656* (STE).

Fig. 4. *Field guide to the Ericas of the Cape Peninsula* (Oliver & Oliver 2000), simplified line drawings by Inge M. Oliver; this opening shows no. 96 *Erica tenuis* Salisb., now *E. oliveranthus* E.C. Nelson & Pirie (Nelson et al 2023: 173–174) and *E. paniculata* L.



Having come into contact with Colonel H. A. Baker by the early 1960s, Ted Oliver became co-author with Baker of *Ericas of South Africa* (Baker & Oliver 1967), illustrated with botanical paintings by the South African botanical artist Irma von Below (1920–1984) and others. Two decades later he started to collaborate with ‘Dolf’ (Adolf Wilhelm Stander) Schumann (1918–2001) and Gerhard P. K. Kirsten (1932–2000) on a profusely illustrated photographic monograph *Ericas of South Africa* (Schumann et al. 1992) in which the amalgamation into the megagenus *Erica* of the last of the minor genera was announced, because they have ‘so much in common ... that their species should also be regarded as ericas [*sic*]’ (Schumann et al. 1992: 244)

Writing for The Heather Society (of which he was an honorary member) in 2000, Ted noted that:

Having begun work on the Ericaceae some 40 years ago as a student, I now find that I am getting to know the group properly, the more so recently because of the rapidly increased knowledge that Inge has also acquired. I am no longer a single person struggling in a “sea of ericas”. There are now two of us to discuss the problems of species de-

limitation, species evolution and the phylogeny, ecology or phytogeography of this fascinating genus. There are quite a number of new species to be described and surprisingly, other un-described species are still being brought in. The biggest problem ahead is that of sub-generic classification. The new techniques of DNA analysis will help to throw some light on this problem, which is now being tackled by a group of international experts with material that I am supplying for them.

Although big plant genera have expanded and contracted over time (Frodin 2004), it is relatively unusual for twentieth-century taxonomists working on morphology alone to change generic delimitations in favour of fewer, larger genera. More often, the emphasis has been placed on morphological differences in particular groups, without necessarily addressing the coherence of groups from which they are split (Humphreys and Linder 2009). Oliver's 'megagenus' concept for *Erica* (Oliver 2012; Oliver & Forshaw 2012) reflected a global understanding of the group that pre-empted subsequent molecular research in which he was instrumental. Phylogenetic trees including more than 40% (Pirie et al. 2011) and 60% (Pirie et al. 2016.) of the species diversity clearly showed that the 'minor genera' are nested within – and indeed scattered across – the redefined mega-genus *Erica*.

One example, *Philippia* Klotzsch, was characterised largely by reduced flowers without brightly coloured corollas or nectaries, but with greatly expanded stigmas (as illustrated in Oliver 1988: 4 & 5), together interpreted as a wind-pollination syndrome (Rebelo et al. 1985). Transition to wind pollination was shown to have occurred several times in *Erica* as then defined (Pirie et al. 2011), with similar characteristics shared by the wind-pollinated 'minor genera' such as *Salaxis* Salisb., *Coccosperma* Klotzsch and *Ericinella* Klotzsch (Oliver 1994, 2000a) and many individual species scattered within *Erica sens. str.*

Blaeria L., had been defined as including those species with four, rather than eight, stamens (Linnaeus 1753, Bentham 1839; Brown 1906; Phillips 1926, 1944; Oliver 1975, 1993b) but this definition was not subsequently applied consistently. Phillips (1944, 1951) included species with four, six or eight stamens in the 'minor genera' *Coccosperma* (4–8 stamens) and *Philippia* (6–8 stamens), but the numbers were also not consistent: four stamens can arise within an individual species usually characterised by having eight (for example, *Erica filiformis* Salisb., *E. blaerioides* E.G.H. Oliv., *E. arborea* L., *E. woodii* Bolus, *E. pleiotricha* S. Moore; Oliver 1993b). As anticipated by Oliver (1993b), neither the 'minor genus' *Blaeria*, nor species of *Erica sens. lat.* possessing only four stamens, proved to represent monophyletic groups, and former *Blaeria* species, such as *E. ericoides* (L.) E.G.H. Oliv., *E. barbigeroide*s E.G.H. Oliv. and *E. russakiana* (Klotzsch ex Walp.) E.G.H. Oliv. from the Cape and the tropical East African species *E. filago* (Alm & T.C.E. Fr.) Beentje and *E. silvatica* (Welw. ex Engl.) Beentje (included in *Erica* by Beentje in 2006, explicitly following Oliver's precedent; Fig. 5), proved to be distantly related.

Fig. 5. Examples of the former genus *Blaeria*. Cape species included in *Erica* by Ted Oliver: A: *E. ericoides*; B. *E. russakiana*; and C: *E. barbigeroides*, and tropical East African species included in *Erica* by Beentje (2006): D: *E. filago* and E: *E. silvatica*. Photos A: MDP; B–E: Berit Gehrke.



With the megagenus concept already firmly embedded, the Olivers set out to revise systematically groups of species following the arrangement in *Flora Capensis* (Guthrie & Bolus 1905, Brown 1906). Two such works were accomplished (Oliver & Oliver 2002; 2005), and many more might have been expected but for the early death of Inge in 2003 (Nelson 2004). Ted continued work on the *Erica* Identification Aid, which includes many of Inge's informal sketches (Oliver et al. 2012; 2024) and provides a route towards species identifications that is also accessible to non-specialists. His support of molecular research from 2008 onwards has included analyses of European species (Kuppler et al. 2015) and improvements to species delimitation in Cape groups (Pirie et al. 2017) as well as broad scale phylogenetic and biogeographic analyses (Pirie et al. 2016; 2019; in prep.) reflecting his broader interests in patterns and origins of Cape plant diversity (Oliver et al. 1983).

Conclusion

By the end of the twentieth century, following almost 250 years of botanical exploration especially in the Cape Floristic Region of South Africa, the number of species of heaths and heathers known to botanists had exceeded eight hundred (Oliver & Oliver 2000) – many hundreds more binomial names had been published (Nelson & Small 2004–2005). By early 2024 the total number of accepted *Erica* species was 851 (Elliot et al. in prep.). Not only is *Erica* confirmed as one of the largest genera of Angiospermae, it is one of the most widely distributed, its species ranging across more than 100 degrees of latitude from northern Norway to the Cape Floristic Province in South Africa, and, straddling the Equator, for almost 90 degrees of longitude from the Azores (+31° W) in the Atlantic Ocean eastwards to islands in the Indian Ocean including Madagascar, the Mascarenes and Mauritius (+57° E) (Oliver 1994a, 2000). At this time, of the currently accepted species, 94 were described and named as new to science by Ted Oliver, and many of these were first collected by him too, while a further 206 are combinations and replacement binomials published under his name, as sole or joint author (Appendix 2). This includes not only Cape diversity, but also taxa from Tropical East Africa and Madagascar (Dorr & Oliver 1999a, 1999b), and the Mascarenes (Oliver 1993a, 2014). Ted and Inge Oliver (pictured in Fig. 6) jointly described and named 16 novel species of *Erica* (Table 1). Inga’s illustrations occurred in many of the papers published since 1974 and (as noted) in the *Erica* Identification Aid (Oliver et al. 2024). A full list of their papers on *Erica* is presented in Appendix 3.

Table 1. New species named by E. G. H. Oliver & I. M. Oliver

<i>Erica amalophylla</i>	E.G.H.Oliv. & I.M.Oliv.	wfo-4000013772
<i>Erica annalis</i>	E.G.H.Oliv. & I.M.Oliv.	wfo-4000013772
<i>Erica cavartica</i>	E.G.H.Oliv. & I.M.Oliv.	wfo-4000013772
<i>Erica ceraria</i>	E.G.H.Oliv. & I.M.Oliv.	wfo-4000013772
<i>Erica croceovirens</i>	E.G.H.Oliv. & I.M.Oliv.	wfo-4000013772
<i>Erica gerhardii</i>	E.G.H.Oliv. & I.M.Oliv.	wfo-4000013772
<i>Erica hebdomalalis</i>	E.G.H.Oliv. & I.M.Oliv.	wfo-4000013772
<i>Erica jananthus</i>	E.G.H.Oliv. & I.M.Oliv.	wfo-4000013772
<i>Erica jugicola</i>	E.G.H.Oliv. & I.M.Oliv.	wfo-4000013772
<i>Erica lithophila</i>	E.G.H.Oliv. & I.M.Oliv.	wfo-4000013772
<i>Erica petrusiana</i>	E.G.H.Oliv. & I.M.Oliv.	wfo-4000013772
<i>Erica prolata</i>	E.G.H.Oliv. & I.M.Oliv.	wfo-4000013772
<i>Erica psittacina</i>	E.G.H.Oliv. & I.M.Oliv.	wfo-4000013772
<i>Erica schelpeorum</i>	E.G.H.Oliv. & I.M.Oliv.	wfo-4000013772
<i>Erica umbratica</i>	E.G.H.Oliv. & I.M.Oliv.	wfo-4000013772
<i>Erica viridimontana</i>	E.G.H.Oliv. & I.M.Oliv.	wfo-4000013772

Fig. 6. Inge M. Oliver (left) and E. G. H. (Ted) Oliver.



No single author has had greater impact on the taxonomy of *Erica*/Ericaceae than Ted Oliver in numbers of new species (as predicted by his mentor, H. A. Baker, in 1962), but this contribution must be interpreted as part of a team effort of two enormously talented botanists. Their combined contribution both of improved knowledge of alpha taxonomy and of broad understanding of the structure of that diversity is fundamental to, and will have a lasting influence on, all future developments in the field.

Acknowledgements

We thank Alan Elliot for help with summarising data from WFO, and Tessa Oliver for consultation.

Authors contributions

ECN: Writing - Original draft, Writing - Review and Editing. MDP: Writing - Review and Editing. DUB: Writing - Review and Editing.

References

- Baker HA (1962) A new species of *Erica* from the Caledon Division. *Journal of South African Botany* 28: 197–199.
[<https://archive.org/details/journalofsouthaf28unse/page/8/mode/2up>]
- Baker HA, Oliver EGH (1967) *Ericas of South Africa*. Purnell, Cape Town.
- Beentje HJ (2006) *Ericaceae*. In: Beentje HJ, Ghazanfar SA (Eds), *Flora of Tropical East Africa*. Royal Botanic Gardens, Kew, 1–29.
- Bentham G (1839) *Ericaceae* (*Ericaceae*). In: De Candolle AP (Ed.), *Prodromus* 7 (2), 612–712.
- Bester SP (2015). *Vaccinium exul* Bolus. [<https://pza.sanbi.org/vaccinium-exul>].
- Bolus H, Guthrie F, Brown NE (1905–1906) Order LXXIX. *Ericaceae*. In Thiselton-Dyer, WT (ed.) *Flora Capensis* vol. 4 section 1, pp. 2–418. Lovell Reeve & Co., London.
- Brown NE (1906) *Philippia*, Klotzsch [to] *Lagenocarpus*, Klotzsch. In Thiselton-Dyer, WT (ed.) *Flora Capensis* vol. 4 section 1, pp. 315–418. Lovell Reeve & Co., London.
- Compton, RH 1935 *Arachnocalyx*, pp 144-146, *Plantae novae Africanae*, series III. *Journal of South African Botany* 1: 111–151
[<https://www.biodiversitylibrary.org/item/231812#page/7/mode/1up>]
- Dorr LJ, Oliver EGH (1999a) A new combination in *Erica* (*Ericaceae*) from São Tomé. *Kew Bulletin* 54: 235–236. Available from: <http://www.jstor.org/stable/4111048>.
- Dorr LJ, Oliver EGH (1999b) New taxa, names, and combinations in *Erica* (*Ericaceae*-*Ericoideae*) from Madagascar and the Comoro Islands. *Adansonia* 21: 75–91. Available from: <https://sciencepress.mnhn.fr/fr/periodiques/adansonia/21/1/nouveaux-taxons-noms-et-combinaisons-dans-le-genre-erica-ericaceae-ericoideae-madagascar-et-aux-comores> (May 12, 2023).
- Elliott A, Bester, SP, Klopper RR, Nelson EC, Pirie MD (in prep) Curating an online checklist for *Erica* L. (*Ericaceae*): contributing to and supporting global conservation through the World Flora Online.
- Frodin DG (2004) History and concepts of big plant genera. *TAXON* 53: 753–776.
<https://doi.org/10.2307/4135449>
- Guthrie F, Bolus H (1905) *Erica*, Linn. In Thiselton-Dyer, WT (ed.) *Flora Capensis* vol. 4 section 1, pp. 4–315. Lovell Reeve & Co., London.
- Humphreys AM, Linder HP (2009) Concept versus data in delimitation of plant genera. *TAXON* 58: 1054–1074. <https://doi.org/10.1002/tax.584002>
- Jarvis C (2007) Order out of chaos. Linnaean plants names and their types. The Linnean Society & the Natural History Museum, London.
- Linnaeus C (1737) *Genera plantarum*. C. Wishoff, Leiden. Available from: <https://doi.org/10.5962/bhl.title.70902>

- Linnaeus C (1753) *Species plantarum*. Vol. 1. L. Salvius, Stockholm. Available from: <https://www.biodiversitylibrary.org/page/26068003>.
- Linnaeus C (1759) *Systema naturae*. L. Salvius, Stockholm
- Linnaeus C (1760) *Plantae rariorae Africanæ*. L. Salvius, Stockholm.
<https://www.biodiversitylibrary.org/page/2031484>
- Linnaeus C (1762) *Species plantarum* ... Editio secunda. Vol. 1. L. Salvius, Stockholm.
- Linnaeus C (1767a) *Systema naturae* ... Editio duodecima. Vol. 3. L. Salvius, Stockholm.
- Linnaeus C (1767b) *Mantissa plantarum*. L. Salvius, Stockholm.
- Linnaeus C (1770) *De Erica*. Edmann, Uppsala.
- Linnaeus C (1771) *Mantissa plantarum altera*. L. Salvius, Stockholm.
- Mugrabi de Kuppler AL, Fagúndez J, Bellstedt DU, Oliver EGH, Léon J, Pirie MD (2015) Testing reticulate versus coalescent origins of *Erica lusitanica* using a species phylogeny of the northern heathers (Ericaceae, Ericaceae). *Molecular Phylogenetics and Evolution* 88: 121–131. <https://doi.org/10.1016/j.ympev.2015.04.005>
- Nelson EC (2004) Inge Oliver (1947–2003). *Heathers: Yearbook of the Heather Society* 1: 55–56. Available from: <https://www.biodiversitylibrary.org/item/148875>.
- Nelson EC (2005) Clarification of a publication date for George Bentham's treatment of Ericaceae (Ericaceae) published in A.-P. de Candolle's *Prodromus* volume 7 part 2 (December 1839). *Archives of natural history* 32: 107. <https://doi.org/10.3366/anh.2005.32.1.106>
- Nelson EC (2011) *Hardy heathers from the northern hemisphere*. Calluna, Daboecia, Erica. A Botanical Magazine monograph. Kew Publishing, Royal Botanic Gardens, Kew.
- Nelson EC, Oliver EGH (2004) Cape heaths in European gardens: the early history of South African *Erica* species in cultivation, their deliberate hybridization and the orthographic bedlam. *Bothalia* 34,2: 127–140. <https://doi.org/10.4102/abc.v34i2.427>
- Nelson EC, Oliver EGH, Pirie MD (2023) *Erica* L. (Ericaceae): homonyms amongst published names for African species and proposed replacement names. *PhytoKeys* 236: 157–178. <https://doi.org/10.3897/phytokeys.236.110498>
- Nelson EC, Pirie MD (2022) Where have all the heathers gone? *Sibbaldia* 21: 75–92. <https://doi.org/10.24823/Sibbaldia.2022.1887>
- Nelson EC, Small DJ (2004–2005) *International register of heather names volume 2 African species, hybrids and cultivars*. 4 parts (part 1 (A–C), 2004; parts 2–4 (D–Z), 2005). The Heather Society, Creeping St Mary.
- Oliver EGH (1964) *Taxonomic studies in the genus Acrostemon Kl. and related genera*. Master of Science thesis. Faculty of Science, University of Cape Town Available from: <http://hdl.handle.net/11427/39023> (accessed January 6, 2024).

- Oliver EGH (1975) Ericaceae. In: Dyer RA (Ed.), The genera of South African flowering plants 1. Botanical Research Institute, Pretoria, 429–439.
- Oliver EGH (1976) Studies in the Ericoideae. II. The new genus *Stokoeanthus*. *Bothalia* 12: 49–52. <https://doi.org/10.4102/abc.v12i1.1370>
- Oliver EGH (1987) Studies in the Ericoideae (Ericaceae). VII. The placing of the genus *Philippia* into synonymy under *Erica*; the southern African species. *South African Journal of Botany* 53: 455–458. [https://doi.org/10.1016/S0254-6299\(16\)31379-5](https://doi.org/10.1016/S0254-6299(16)31379-5)
- Oliver EGH (1988) Studies in the Ericoideae (Ericaceae). VI. The generic relationship between *Erica* and *Philippia* in southern Africa. *Bothalia* 18: 1–10. <https://doi.org/10.4102/abc.v18i1.975>
- Oliver EGH (1993a) Studies in the Ericoideae (Ericaceae). X. Nomenclatural Changes for the Flore des Mascareignes Region. *Kew Bulletin* 48: 767–769. <https://doi.org/10.2307/4118855>
- Oliver EGH (1993b) Studies in the Ericoideae (Ericaceae). XI. The generic relationship between *Erica* and *Blaeria*. *Kew Bulletin* 48: 771–780. <https://doi.org/10.2307/4118856>
- Oliver EGH (1993c) Studies in the Ericoideae (Ericaceae). XII. The placing of the genus *Blaeria* into synonymy under *Erica*; nomenclatural and taxonomic changes for the southern African region. *Bothalia* 23: 1–7. <https://doi.org/10.4102/abc.v23i1.779>
- Oliver EGH (1994) Studies in the Ericoideae (Ericaceae). XV. The generic relationship between *Erica* and *Ericinella*. *Bothalia* 24: 121–126. <https://doi.org/10.4102/abc.v24i2.761>
- Oliver EGH (1996) The position of *Bruckenthalia* versus *Erica*. *Yearbook of the Heather Society*: 6. Available from: <https://www.biodiversitylibrary.org/page/43695628>.
- Oliver EGH (1999) Systematic studies in the Tribe Ericaceae (Ericaceae–Ericoideae). Doctoral thesis. University of Cape Town.
- Oliver EGH (2000a) Systematics of Ericaceae (Ericaceae: Ericoideae) species with indehiscent and partially dehiscent fruits. *Contributions from the Bolus Herbarium* No. 19.
- Oliver EGH (2000b) An obsession with *Erica*. *Yearbook of The Heather Society* 2000: 52–56. <https://www.biodiversitylibrary.org/page/43695473>
- Oliver EGH (2004) *Erica*. In: Kubitzki K (Ed.), The families and genera of vascular plants, volume 6. Flowering Plants. Dicotyledons: Celastrales, Oxalidales, Rosales, Cornales, Ericales. p. 173. Springer Berlin Heidelberg, Berlin, Heidelberg. https://doi.org/10.1007/978-3-662-07257-8_19
- Oliver EGH (2007) Linnaeus' Cape species of *Erica*. *Heathers* 4: 5–10. Available from: <https://www.biodiversitylibrary.org/page/43784094>.
- Oliver EGH (2012) Ericaceae. In: Manning J, Goldblatt P (Eds.) *Plants of the Greater Cape Floristic Region. 1: The core Cape Flora. Strelitzia* 29: 482–511.
- Oliver EGH (2014) *Ericas* in Mauritius, home of the dodo. *Heathers* 11: 38–42.

<https://doi.org/10.5281/zenodo.10361982>

Oliver EGH, Forshaw N (2012) Genus *Erica* An Identification Aid Version 3.00. Contributions from the Bolus Herbarium 22.

Oliver EGH, Forshaw N, Oliver IM, Volk F, Schumann AWS, Dorr LJ, Hoekstra RD, Musker SD, Nürk NM, Pirie M, Rebelo AG (2024) Genus *Erica*: An Identification Aid Version 4.00. ARPHA Preprints 5: e117930. <https://doi.org/10.3897/arphapreprints.e117930>

Oliver EGH, Linder HP, Rourke JP (1983) Geographical distribution of present-day Cape taxa and their phytogeographical significance. *Bothalia* 14: 427–440. <https://doi.org/10.4102/abc.v14i3/4.1189>

Oliver EGH, Oliver IM (1991) Studies in the Ericoideae (Ericaceae). VIII. New species in *Erica*, section *Pseuderemia*, from southern Africa. *Bothalia* 21: 137–142. <https://doi.org/10.4102/abc.v21i2.873>

Oliver EGH, Oliver IM (2000a) Ericaceae. In: Goldblatt P, Manning J (Eds), Cape plants: a conspectus of the Cape flora of South Africa. *Strelitzia* 9. National Botanical Institute, 423–452. Available from: <https://www.cabdirect.org/cabdirect/abstract/20013066367> (October 4, 2023).

Oliver I[M], Oliver T [EGH] (2000b) Field guide to the ericas of the Cape Peninsula. Protea Atlas Project, National Botanical Institute.

Oliver EGH, Oliver IM (2002) The genus *Erica* (Ericaceae) in southern Africa: taxonomic notes 1. *Bothalia* 32: 37–61. <https://doi.org/10.4102/abc.v32i1.461>

Oliver EGH, Oliver IM (2005) The genus *Erica* (Ericaceae) in southern Africa: taxonomic notes 2. *Bothalia* 35,2: 121–148. <https://doi.org/10.4102/abc.v35i2.388>

Pirie MD, Oliver EGH, Bellstedt DU (2011) A densely sampled ITS phylogeny of the Cape flagship genus *Erica* L. suggests numerous shifts in floral macro-morphology. *Molecular Phylogenetics and Evolution* 61: 593–601. <https://doi.org/10.1016/j.ympev.2011.06.007>

Pirie MD, Oliver EGH, Mugrabi de Kuppler A, Gehrke B, Le Maitre NC, Kandziora M, Bellstedt DU (2016) The biodiversity hotspot as evolutionary hot-bed: spectacular radiation of *Erica* in the Cape Floristic Region. *BMC Evolutionary Biology* 16: 190. <https://doi.org/10.1186/s12862-016-0764-3>

Pirie MD, Oliver EGH, Gehrke B, Heringer L, Mugrabi de Kuppler A, Le Maitre NC, Bellstedt DU (2017) Underestimated regional species diversity in the Cape Floristic Region revealed by phylogenetic analysis of the *Erica abietina*/*E. viscaria* clade (Ericaceae). *Botanical Journal of the Linnean Society* 184: 185–203. <https://doi.org/10.1093/botlinnean/box021>

Pirie MD, Kandziora M, Nürk NM, Le Maitre NC, Mugrabi de Kuppler A, Gehrke B, Oliver EGH, Bellstedt DU (2019) Leaps and bounds: geographical and ecological distance constrained the colonisation of the Afrotropical by *Erica*. *BMC Evolutionary Biology* 19: 222. <https://doi.org/10.1186/s12862-019-1545-6>

Phillips EP (1926) The genera of South African flowering plants. Botanical Survey Memoir no. 10. <http://archive.org/details/generaofsouthafr00phil> (January 17, 2024).

Phillips EP (1944) Notes on the minor genera of the Ericaceae. Journal of South African Botany 10: 69–73. <https://archive.org/details/journalofsouthaf10unse/page/68/mode/2up> (February 13, 2024)

Phillips EP (1951) The genera of South African flowering plants. 2nd ed. Botanical Survey Memoir 25, 923 pp. https://archive.org/details/generaofsouthafr00phil_0/page/561/mode/1up (February 13, 2024).

Rebello AG, Siegfried WR, Oliver EGH (1985) Pollination syndromes of *Erica* species in the south-western Cape. South African Journal of Botany 51: 270–280. [https://doi.org/10.1016/S0254-6299\(16\)31657-X](https://doi.org/10.1016/S0254-6299(16)31657-X)

Schumann D, Kirsten G, Oliver EGH (1992) *Ericas of South Africa*. Fernwood Press, Vlaeberg, South Africa, 272 pp.

Appendices

Appendix 1: Generic description of *Erica* by E. G. H. Oliver (2000a; from Nelson 2011: 100)

Appendix 2: A complete list of *Erica s.l.* taxa at all ranks authored by E.G.H. Oliver and I.M. Oliver, including taxa new to science, replacement names, and new combinations, with synonymy and bibliographic data, derived from the World Flora Online (Elliot et al., in prep.).

Appendix 3: A complete list of *Erica* related publications authored by E.G.H. Oliver and I.M. Oliver.