

PREPRINT

Posted on 04/11/2020

DOI: <https://doi.org/10.3897/arphapreprints.e60028>

A global food plant dataset for wild silkmoths and hawkmoths, and its use in documenting polyphagy of their caterpillars (Lepidoptera: Bombycoidea: Saturniidae, Sphingidae)

 Liliana Ballesteros Mejia, Pierre Arnal, Winnie Hallwachs, Jean Haxaire, Daniel Janzen, Ian Kitching,  Rodolphe Rougerie

Not peer-reviewed, not copy-edited manuscript.

A global food plant dataset for wild silkmoths and hawkmoths, and its use in documenting polyphagy of their caterpillars (Lepidoptera: Bombycoidea: Saturniidae, Sphingidae)

Liliana Ballesteros Mejia^{‡,§,||}, Pierre Arnat[‡], Winnie Hallwachs[¶], Jean Haxaire^{#,□}, Daniel Janzen[«], Ian J. Kitching[»], Rodolphe Rougerie[‡]

‡ Institut de Systématique, Evolution, Biodiversité (ISYEB), Muséum national d'Histoire naturelle, CNRS, Sorbonne Université, EPHE, Université des Antilles (1st authorship shared between the first two authors), Paris, France

§ CESAB, Centre de Synthèse et d'Analyse sur la Biodiversité, Montpellier, France

| Ecologie, Systématique and Evolution, Université Paris-Sud, CNRS, AgroParisTech, Université Paris-Saclay, Orsay, France

¶ Department of Biology, University of Pennsylvania, Philadelphia, United States of America

Correspondent of Muséum national d'Histoire naturelle, Paris, France

□ Associate researcher of Insectarium de Montréal, Quebec, Canada

« University of Pennsylvania, Philadelphia, United States of America

» Department of Life Sciences, Natural History Museum, London, United Kingdom

Corresponding author: Rodolphe Rougerie (rodolphe.rougerie@mnhn.fr)

Abstract

Background

Herbivorous insects represent a major fraction of global biodiversity and the relationships they have established with their food plants range from strict specialists to broad generalists. Our knowledge of these relationships is of primary importance to basic (e.g. the study of insect ecology and evolution) and applied biology (e.g. monitoring of pest or invasive species), and yet remains very fragmentary and understudied. In Lepidoptera, caterpillars of families Saturniidae and Sphingidae are rather well known and considered to have adopted contrasting preferences in their use of food plants. The former are regarded as being rather generalist feeders, whereas the latter are more specialist.

New information

To assemble and synthesize the vast amount of existing data on food plants of Lepidoptera families Saturniidae and Sphingidae, we combined three major existing databases to produce a dataset collating more than 26,000 records for 1256 species (25% of all species) in 121 (67%) and 167 (81%) genera of Saturniidae and Sphingidae, respectively. This dataset is used here to document the level of polyphagy of each of these genera using

summary statistics as well as the calculation of a polyphagy score derived from the analysis of Phylogenetic Diversity of the food plants used by the species in each genus.

Keywords

Lepidoptera, food plant, ecology, life-history traits, caterpillar, polyphagy

Introduction

Herbivorous insects represent a major fraction of global biodiversity (Fiedler 1998) and are central to studies of numerous and diverse ecological and evolutionary processes, such as resource specialization (Devictor et al. 2008), coevolution (Thompson 1988) and food web dynamics (Vidal and Murphy 2017). Elucidating the degree of food plant-insect specificity helps understand community assembly, ecosystem dynamics and latitudinal gradients of species richness (Ødegaard 2006). Moreover, insect-plant interactions are central to the understanding of niche breadth and they play a key role in mediating competition that structures communities, and backdrop the human view of entire networks of interacting species (Devictor et al. 2008, Forister et al. 2014). The different levels of specialization observed in phytophagous insects, from strict specialists to highly generalist species, are traits that are also considered as possibly important drivers of speciation or adaptive radiation (Janz and Nylin 2008, Jousset and Elias 2019, Wang et al. 2017).

The Lepidoptera families Saturniidae (wild silkmoths) and Sphingidae (hawkmoths, sphinx moths) are among the best-known insect families worldwide, both taxonomically and biologically, and they are generally characterized by being large-bodied moths (Janzen 1984a). A recently published taxonomic checklist (Kitching et al. 2018) revealed a combined species richness of around 5000 species globally. These two families exhibit contrasting life-history strategies both as adults - Sphingidae (feeding, long-lived adults) and Saturniidae (non-feeding, short-lived adults) (Janzen 1984a) - and as caterpillars - Sphingidae (fast growing, many toxic plant specialists) and Saturniidae (slow-growing, many tannin and resin-rich plant specialists) (Janzen 1981, Janzen 1984a). In the Neotropics sphingid caterpillars seem to specialize on only a relatively small number of plant families, feeding on both young and old, relatively tender leaves that contain low molecular weight toxic compounds, whereas saturniid caterpillars feed on tougher, as well as younger, leaves of an often wide range of plant families that contain high levels of large polymeric molecules (tannins, resins) that interfere with digestion (Janzen 1981). Consequently, sphingid caterpillars digest more nutrients per bite and need less time to reach a given full size than do saturniids (Bernays and Janzen 1988).

A massive amount of data is available on the larval food plants in the wild of the two families, both in the literature and in institutional and personal databases. For the Lepidoptera as a whole, the HOSTS database (Robinson et al. 2010) comprises the most comprehensive collation of information about what caterpillars overall are believed to eat. It contains some 180,000 records for about 22,000 Lepidoptera species extracted from 1600

documents (Robinson et al. 2010). Although HOSTS has not been updated for almost a decade, the subset of records for the superfamily Bombycoidea has been independently maintained and added to by IJK and this updated version is used here. Another spectacular effort towards gathering food plant data for Lepidoptera is the inventory of caterpillars in the Area de Conservacion Guanacaste (ACG) in northwestern Costa Rica (Janzen and Hallwachs 2016, Janzen and Hallwachs 2020). It comprises ~70,000 records of reared wild-caught larvae of Saturniidae and Sphingidae linked to their DNA barcodes. Besides these two main public data repositories, one of the authors (JH) has built his own personal database for Sphingidae over 20 years, compiling records from the literature, web resources, personal field observations and communications from collaborators. In addition, food plant information is also scattered across the published literature, including a few more recent food plant catalogues, such as in Stone (1994), Santin (2004), Meister (2011), but also webpages and personal databases, all of which makes the process of collating and resolving the information very difficult and time consuming.

All three databases cited above are and remain independently maintained and updated. Here we publish a single dataset resulting from their combination. Our aim is to make this massive amount of information available as a single dataset that allows its use for ecological and evolutionary analyses. In particular, we want to investigate the role of food plant use in the evolution of the two families (Arnal et al., in prep.), especially with respect to the degree of polyphagy, defined as the plasticity in the use of different food plants for caterpillars to complete their development. We provide further details about the contents of this dataset in the following sections, as well as a number of caveats to avoid incorrect interpretation and use of these data. In addition to variables summarizing the level of polyphagy of the caterpillars of sphingid and saturniid moths, we also provide a polyphagy score based on a calculation of Phylogenetic Diversity (Faith 1992) of the food plant families used by the species included in the database.

General description

Purpose: The food plant dataset

This dataset (Suppl. material 1) is a synthesis of current knowledge regarding the food plants eaten by the caterpillars of two families of Lepidoptera (Saturniidae and Sphingidae). It aims to capture the state of knowledge at the time of assembly of the dataset so that it can be used to investigate the role of food plants use breadth in the spatial and temporal evolution of both families (Arnal et al., in prep.).

This dataset of larval food plant records for sphingids and saturniids worldwide is the result of the integration, with significant data reconciliation and standardization, of these three largely independent data sources:

1) Information for Sphingidae and Saturniidae embedded in the HOSTS database (Robinson et al. 2010); as further added to and refined by IJK, downloaded on March 2nd, 2018) (hereafter HOSTS);

2) An inventory of the caterpillars, their food plants and parasitoids of Area de Conservacion Guanacaste (ACG, Janzen DH, downloaded on July 16th 2018 for Saturniidae and July 18th 2018 for Sphingidae) (hereafter DHJ);

3) The personal database of Jean Haxaire (Associate Researcher to MNHN, imported on July 17th 2018) (hereafter JH).

A “record” refers to a unique combination of caterpillar species, plant species and source. Records in the dataset resulting from rearing experiments in captivity or from introduced plant species are listed separately as they often do not represent natural insect-plant associations. Redundancy (duplication) of records among the three databases following their combination was not a concern for our research objectives; the dataset should be treated as qualitative, and the frequency of records ignored (see list of points in next section).

A total of 25,937 records was compiled from the three databases in a single dataset given as Suppl. material 1. Table 1 below provides details of the number of records contributed by each of the independent databases. We followed the plant taxonomy of the International Plant Names Index (IPNI; <https://www.ipni.org>) and the latest moth taxonomy (Kitching et al. 2018), though both do not coincide with some of the names used by all three sources (see ‘call for caution’ below).

This compilation provides information for 137 genera and 757 species of Saturniidae and 166 genera and 725 species of Sphingidae.

As an example of the uses of this dataset, we report basic polyphagy variables as well as a polyphagy score based on Phylogenetic Diversity (PD, Faith 1992) of the food plants used by the caterpillars of saturniid and sphingid moths. Using a recent dated angiosperm phylogeny (Magallón et al. 2015) we measured the PD score, i.e. the total length of all phylogenetic tree branches connecting the different families of plants eaten by a given moth species *in natura*, using the *pd* function of the *picante* R package (Kembel et al. 2010). The species scores were then averaged within each genus to get genus scores in Suppl. material 2. Note that gymnosperm records were excluded from our calculations of PD scores to avoid bias caused by the considerable phylogenetic distance between angiosperms and gymnosperms.

The genus-level polyphagy variables and the polyphagy scores of Saturniidae and Sphingidae genera are provided as Suppl. material 2.

Additional information: Calls for caution:

1. The correctness of food plant identifications in databases and in the literature should be treated with considerable caution, as they were largely made by non-

- botanists; food plant names used are also subject to taxonomic and nomenclatural uncertainty, and their correctness and validity may be considered equivocal in some cases.
2. The previous point also applies to moth names, especially when considering species-level identifications. These may be incorrect or outdated. For example, more than 1500 new species have been described within family Saturniidae in the past decade (Kitching et al. 2018), largely with the support of DNA barcoding analyses. Thus, food plant records may not account for recently split complexes of cryptic species, members of which may have quite different natural histories (e.g. Janzen (2012)).
 3. The food plant dataset is derived from known food plant records at the time of its compilation; as such it represents a snapshot of the knowledge at that time and it may differ from the data compiled in the original sources and then updated independently (e.g. new records and/or corrections (e.g. identification errors or synonymies of the moth/caterpillar, or the plant, or both)).
 4. All records are meant to represent actual instances in which caterpillars were found feeding and developing on the food plant. Records in the DHJ database all result from rearing trials of caterpillars found in the field on the food plant in question, and in many cases, identification of the caterpillar was confirmed through DNA barcoding of the resultant adult moths. A few records recognized as questionable (e.g. inconsistent locality/identification data) in the HOSTS and JH databases were filtered out and are not included in the present combined dataset.
 5. The food plant dataset does not account for the frequency of use of a given food plant among other plants also listed for the same species of moth. The DHJ database, because it is based on individual specimen records, does include quantitative data; however, this information is not incorporated into the combined dataset, although it could bring additional information on local food plant preferences of species and populations. We note that this information would nevertheless be very difficult to analyze and interpret as it is conditional upon the local availability of food plants, as well as possibly seasonal conditions, local variations through time and difficulty of collecting.
 6. The previous point also brings a note of caution in that polyphagy, as calculated here from the data available for a given species, may not be translatable to the population or site level, and vice versa. A species may have populations in which some caterpillars have a lower level of polyphagy than others, at least in part because the food plants that could be eaten do not occur in that ecosystem and because many species arrive by ecological fitting rather than *in situ* evolution (Janzen 1985a). This is especially the case with species following expanding frontier agriculture into new ecosystems, or following contemporary climate changes.
 7. Strictly speaking, we define polyphagy as the capacity of a given individual caterpillar to feed and develop (through its complete life cycle) on different food plants. This can only be approximated by considering sibling individuals (as is sometimes the case in DHJ database), individuals from the same population, or, ultimately from the same species or higher taxonomic categories. We thus

- acknowledge that the scores of polyphagy at species and genus level should be recognized as human abstractions.
8. Polyphagy is constrained *in situ* by the local availability of food plants - an individual caterpillar cannot be polyphagous on species of plants that are not present.
 9. Here we approximated polyphagy scores at species level for saturniid and sphingid moths, and we assume that they represent valuable information about the level of plasticity of individuals of the populations of a species to use different food plants. These scores were then used to calculate polyphagy scores at the genus level. Generic level of polyphagy is a human abstraction, but it is seen as relevant information to understand the past diversification dynamics. Plasticity in the use of food plants may have favored or impeded geographical dispersal, and may have mitigated speciation or extinction processes, or influenced species natural histories in many other ways (Janzen 1985b).
 10. We acknowledge that the polyphagy level derived from caterpillar plant feeding records approximates, but may not reflect precisely, the plasticity in oviposition site selection by female moths (see for instance Janzen 1984b). Indeed, caterpillars may be driven by starvation to feed on a different plant after consuming all leaves of the plant they started to develop on and which had been selected for oviposition by the female.

Geographic coverage

Description: The present dataset combines food plant records for saturniid and sphingid species worldwide.

Taxonomic coverage

Taxa included:

Rank	Scientific Name	Common Name
family	Saturniidae	Wild silkmoths
family	Sphingidae	Hawkmoths

Usage licence

Usage licence: Open Data Commons Attribution License

Data resources

Data package title: Global food plant dataset and polyphagy scores for Sphingidae and Saturniidae

Number of data sets: 2

Data set name: Global food plant dataset for Saturniidae and Sphingidae species

Data format: Excel data spreadsheet

Description: Suppl. material 1

Column label	Column description
Family	Taxonomic family of the moth genus/species
Subfamily	Taxonomic subfamily of the moth genus/species
Tribe	Taxonomic tribe of the moth genus/species
Moth_Genus_name	Genus name
Moth_Species_Name	Species name
Number_PlantGenus	Total number of plant genera known to be eaten in natural environment by caterpillars of this species of moth
Plant_GenusNames	Names of plant genera known to be eaten in natural environment by caterpillars of this species of moth
Number_PlantSpecies	Total number of plant species known to be eaten in natural environments by caterpillars of this species of moth
Plant_SpeciesNames	Names of plant species known to be eaten in natural environments by caterpillars of this species of moth
Number_PlantFamily	Total number of plant families known to be eaten in natural environments by caterpillars of this species of moth
Plant_FamilyNames	Names of plant families known to be eaten in natural environments by caterpillars of this species of moth
Number_PlantOrders	Total number of plant orders known to be eaten in natural environments by caterpillars of this species of moth
Plant_OrderNames	Names of plant orders known to be eaten in natural environments by caterpillars of this species of moth
Number_PlantGenus_Capt	Total number of plant genera known to be eaten in captivity by caterpillars of this species of moth
Plant_GenusNames_Capt	Names of plant genera known to be eaten in captivity by caterpillars of this species of moth
Number_PlantSpecies_Capt	Total number of plant species known to be eaten in captivity by caterpillars of this species of moth
Plant_SpeciesNames_Capt	Names of plant species known to be eaten in captivity by caterpillars of this species of moth

Number_PlantFamily_Capt	Total number of plant families known to be eaten in captivity by caterpillars of this species of moth
Plant_FamilyNames_Capt	Names of plant families known to be eaten in captivity by caterpillars of this species of moth
Number_PlantOrders_Capt	Total number of plant orders known to be eaten in captivity by caterpillars of this species of moth
Plant_OrderNames_Capt	Names of plant orders known to be eaten in captivity by caterpillars of this species of moth

Data set name: Polyphagy variables and score for Saturniidae and Sphingidae

Download URL:

Data format: Excel data spreadsheet

Description: Suppl. material 2

Column label	Column description
Family	Taxonomic family of the moth genus
Subfamily	Taxonomic subfamily of the moth genus
Tribe	Taxonomic tribe of the moth genus
Moth_Genus_Name	Genus name
NumberSampledMothSpecies	Number of moth species within the genus that have food plant information available
TotalMothSpecies	Total number of moth species within the genus
TotalNumberGenus	Total number of plant genera known to be eaten in natural environment by caterpillars of this genus of moth
AverageNumberGenus	The average number of plant genera known to be eaten in natural environments by species within this genus of moth
TotalNumberFamilies	Total number of plant families known to be eaten in natural environments by caterpillars of this genus of moth
AverageNumberFamilies	The average number of plant families known to be eaten in natural environments by species within this genus of moth
TotalNumberOrders	Total number of plant orders known to be eaten in natural environments by caterpillars of this genus of moth
AverageNumberOrders	The average number of plant orders known to be eaten in natural environments by species within this genus of moth

PD_score	Phylogenetic score based on the total branch length in the phylogenetic tree connecting the different families of plant eaten by a given moth species and summarized by genus. NOTE: Only computed from records of plant species eaten in natural environments; records on Gymnosperms were discarded prior calculation (see text).
----------	---

Acknowledgements

- This work is a product of the ACTIAS group funded by the synthesis center CESAB of the French Foundation for Research on Biodiversity (FRB; www.fondationbiodiversite.fr).
- PA and RR are supported by ANR grant SPHINX ANR-16-CE02-0011-01.
- We thank the entire ACG parataxonomist team (see Janzen and Hallwachs, 2020) for finding, rearing and processing the specimens described and discussed here in database DHJ, and we thank the Centre for Biodiversity Genomics at the University of Guelph for DNA barcoding them and otherwise managing their molecular data; Area de Conservacion Guanacaste/MINAE and donors to the Guanacaste Dry Forest Conservation Fund for preserving the ACG rain forest in which they live; the Systematic Entomology Laboratory of the USDA and US National Museum/ Smithsonian Institution for receiving and permanently housing the specimens and providing the human resources to carry out this taxonomic work, and finally, many dozens of members of the taxasphere for applying names to both the moths and their food plants.
- IJK would like to thank the Trustees of the Loke Wan Tho Memorial Foundation for their generous support of the HOSTS project and his colleagues at the NHM who undertook the original HOSTS project: Phillip Ackery, George Beccaloni, Luis Hernández, Adrian Hine, Sven Loburg, Mike Lowndes and most of all, the late Gaden Robinson, whose dedication saw the project to completion. He is also extremely grateful to the many people who contributed their own rearing records of Lepidoptera or personal accumulations of data for inclusion in the HOSTS database, particularly Mike Bigger (UK), John W. Brown (USA), Chris Conlan (USA), Rob Ferber (USA), Konrad Fiedler (Germany), Jeremy Holloway (UK), Frank Hsu (USA), Jurie Intachat (Malaysia), Alec McClay (Canada), Bill Palmer (Australia), Pierre Plauzoles (USA) and the generous individuals who contributed rearing records through the WorldWideWeb and who are known to us only as an email address. IJK is particularly grateful to Julian Donahue and the Los Angeles County Museum of Natural History for allowing us to include data into HOSTS on Microlepidoptera from the card catalogue prepared by the late J.A. Comstock and C. Henne, and for access to manuscript records by Noel McFarland. Full acknowledgements for the HOSTS database can be found at <https://www.nhm.ac.uk/our-science/data/hostplants/#9>.

Author contributions

PA, LBM, IJK and RR designed the study and organized the assembly of the dataset. JH, IJK, WH and DHJ compiled the three databases; LBM carried out their combination and computed summary statistics of polyphagy levels. PA computed the calculation of polyphagy scores.

LBM wrote the first draft of the manuscript, then all authors contributed to its redaction and to the edition of its final version.

References

- Bernays EA, Janzen DH (1988) Saturniid and sphingid Caterpillars: Two ways to eat Leaves. *Ecology* 69 (4): 1153-1160. <https://doi.org/10.2307/1941269>
- Devictor V, Julliard R, Jiguet F (2008) Distribution of specialist and generalist species along spatial gradients of habitat disturbance and fragmentation. *Oikos* 117 (4): 507-514. <https://doi.org/10.1111/j.0030-1299.2008.16215.x>
- Faith D (1992) Conservation evaluation and phylogenetic diversity. *Biological Conservation* 61 (1): 1-10. [https://doi.org/10.1016/0006-3207\(92\)91201-3](https://doi.org/10.1016/0006-3207(92)91201-3)
- Fiedler K (1998) Diet breadth and host plant diversity of tropical- vs. temperate-zone herbivores: South-East Asian and West Palaearctic butterflies as a case study. *Ecological Entomology* 23 (3): 285-297. <https://doi.org/10.1046/j.1365-2311.1998.00132.x>
- Forister M, Novotny V, Panorska A, Baje L, Basset Y, Butterill P, Cizek L, Coley P, Dem F, Diniz I, Drozd P, Fox M, Glassmire A, Hazen R, Hrcek J, Jahner J, Kaman O, Kozubowski T, Kursar T, Lewis O, Lill J, Marquis R, Miller S, Morais H, Murakami M, Nickel H, Pardikes N, Ricklefs R, Singer M, Smilanich A, Stireman J, Villamarin-Cortez S, Vodka S, Volf M, Wagner D, Walla T, Weiblen G, Dyer L (2014) The global distribution of diet breadth in insect herbivores. *Proceedings of the National Academy of Sciences* 112 (2): 442-447. <https://doi.org/10.1073/pnas.1423042112>
- Janzen DH (1981) Patterns of herbivory in a tropical deciduous forest. *Biotropica* 13 (4). <https://doi.org/10.2307/2387805>
- Janzen DH (1984a) Two ways to be a tropical big moth: Santa Rosa saturniids and sphingids. *Oxford Surveys in Evolutionary Biology* 1: 85-140.
- Janzen DH (1984b) Natural history of *Hylesia lineata* (Saturniidae: Hemileucinae) in Santa Rosa National Park, Costa Rica. *Journal of the Kansas Entomological Society* 57: 490-514.
- Janzen DH (1985a) On ecological fitting. *Oikos* 45: 308-310.
- Janzen DH (1985b) A host plant is more than its chemistry. *Illinois Natural History Bulletin* 33: 141-174.
- Janzen DH, et al. (2012) What happens to the traditional taxonomy when a well-known tropical saturniid moth fauna is DNA barcoded? *Invertebrate Systematics* 26 (6): 478-505. <https://doi.org/https://doi.org/10.1071/IS12038>

- Janzen DH, Hallwachs W (2016) DNA barcoding the Lepidoptera inventory of a large complex tropical conserved wildland, Area de Conservacion Guanacaste, northwestern Costa Rica. *Genome* 59 (9): 641-660. <https://doi.org/10.1139/gen-2016-0005>
- Janzen DH, Hallwachs W (2020) Caterpillars, pupae, butterflies & moths of ACG, Guanacaste, Costa Rica. <http://Janzen.sas.upenn.edu/caterpillars/database.lasso>. Accessed on: 2020-9-18.
- Janz N, Nylin S (2008) The oscillation hypothesis of host-plant range and speciation. In: Tilman D (Ed.) *The evolutionary biology of herbivorous insects: specialization, speciation and radiation*. University of California Press, 203-215 pp.
- Jousselin E, Elias M (2019) Testing host-plant driven speciation in phytophagous Insects: A phylogenetic perspective. *BioArXiv*. 2019020215 <https://doi.org/10.20944/preprints201902.0215.v1>
- Kembel SW, Cowan PD, Helmus MR, Cornwell WK, Morlon H, Ackerly DD, Blomberg SP, Webb CO (2010) Picante: R tools for integrating phylogenies and ecology. *Bioinformatics* 26 (11): 1463-1464.
- Kitching I, Rougerie R, Zwick A, Hamilton C, St Laurent R, Naumann S, Ballesteros-Mejia L, Kawahara A (2018) A global checklist of the Bombycoidea (Insecta: Lepidoptera). *Biodiversity Data Journal* 6: e22236. <https://doi.org/10.3897/bdj.6.e22236>
- Magallón S, Gómez-Acevedo S, Sánchez-Reyes L, Hernández-Hernández T (2015) A metacalibrated time-tree documents the early rise of flowering plant phylogenetic diversity. *New Phytologist* 207 (2): 437-453. <https://doi.org/10.1111/nph.13264>
- Meister F (2011) *Die Zucht von tropischen Wilden Seidenspinnern (Lepidoptera: Saturniidae)*. [A guide to the breeding of tropical silk moths]. Verlag Dr. Friedrich Pfeil, Munchen, 220 pp.
- Ødegaard F (2006) Host specificity, alpha- and beta-diversity of phytophagous beetles in two tropical forests in Panama. *Biodiversity and Conservation* 15 (1): 83-105. <https://doi.org/10.1007/s10531-004-3106-5>
- Robinson GS, Ackery PR, Kitching IJ, Beccaloni GW, Hernandez LM (2010) A database of the World's Lepidopteran hostplants. Natural History Museum, London. <http://www.nhm.ac.uk/hosts>. Accessed on: 2020-10-23.
- Santin A (2004) *Répertoire des Plantes-Hôtes de Substitution des chenilles du monde*. Éléments recueillis dans des publications entomologiques et résultats d'expérimentations personnelles. Deuxième édition. Office pour l'information éco-entomologique, Guyancourt, 1228 pp.
- Stone SE (1994) Foodplants of World Saturniidae. *Memoirs of the Lepidopterists' Society* 4: 1-186. URL: <https://www.lepsoc.org/catalog/memoirs>
- Thompson J (1988) Coevolution and alternative hypotheses on insect/plant interactions. *Ecology* 69 (4): 893-895. <https://doi.org/10.2307/1941238>
- Vidal M, Murphy S (2017) Bottom-up vs. top-down effects on terrestrial insect herbivores: a meta-analysis. *Ecology Letters* 21 (1): 138-150. <https://doi.org/10.1111/ele.12874>
- Wang H, Holloway J, Janz N, Braga M, Wahlberg N, Wang M, Nylin S (2017) Polyphagy and diversification in tussock moths: Support for the oscillation hypothesis from extreme generalists. *Ecology and Evolution* 7 (19): 7975-7986. <https://doi.org/10.1002/ece3.3350>

Table 1.

General overview of the contribution of each database to our dataset (Suppl. material 1).

Database	Family	Number of records	Geographical coverage
HOSTS	Saturniidae	10586	Worldwide
	Sphingidae	10528	Worldwide
DHJ	Saturniidae	2297	Local in three adjoining ecosystems
	Sphingidae	1322	Local in three adjoining ecosystems
JH	Sphingidae	2401	Worldwide

Supplementary materials

Suppl. material 1: Food plant dataset for worldwide saturniid and sphingid moths (Lepidoptera, Saturniidae, Sphingidae)

Authors: Kitching, I.J., Janzen, D.H.J., Hallwachs, W, Haxaire, J

Data type: food plant and moths association records

Brief description: This dataset lists food plants known to be fed on by caterpillars of saturniid and sphingid moths, worldwide. It includes both wild and captive records (listed separately).

[Download file](#) (320.46 kb)

Suppl. material 2: Polyphagy variables and score for Saturniidae and Sphingidae

Authors: Ballesteros-Mejia, L., Arnal, P., Kitching, I.J., Rougerie, R.

Data type: Summary variables, polyphagy score

Brief description: This table reports summary variables and score of polyphagy for Saturniidae and Sphingidae records of our dataset. Note that gymnosperm records were excluded from our calculations of PD scores to avoid bias caused by the considerable phylogenetic distance between angiosperms and gymnosperms. Polyphagy scores were not calculated for genera feeding on Gymnosperms only .

[Download file](#) (31.44 kb)