

PREPRINT

Author-formatted, not peer-reviewed document posted on 29/06/2021

DOI: <https://doi.org/10.3897/raphapreprints.e70782>

Gene Sequence Analysis of Mitochondrial COI in Genus *Saperda* (Coleoptera: Cerambycidae: Lamiinae)

 Wei XU, Qinhua Gan, Jian Pu, Yingwen Pan, Bo Cai, Su Ao, Rui Meng, Li Chen

Disclaimer on biological nomenclature and use of preprints

The preprints are preliminary versions of works accessible electronically in advance of publication of the final version. They are not issued for purposes of botanical, mycological or zoological nomenclature and **are not effectively/validly published in the meaning of the Codes**. Therefore, nomenclatural novelties (new names) or other nomenclatural acts (designations of type, choices of priority between names, choices between orthographic variants, or choices of gender of names) **should NOT be posted in preprints**. The following provisions in the Codes of Nomenclature define their status:

International Code of Nomenclature for algae, fungi, and plants (ICNafp)

Article 30.2: "An electronic publication is not effectively published if there is evidence within or associated with the publication that its content is merely preliminary and was, or is to be, replaced by content that the publisher considers final, in which case only the version with that final content is effectively published." In order to be validly published, a nomenclatural novelty must be effectively published (Art. 32.1(a)); in order to take effect, other nomenclatural acts must be effectively published (Art. 7.10, 11.5, 53.5, 61.3, and 62.3).

International Code of Zoological Nomenclature (ICZN)

Article: 21.8.3: "Some works are accessible online in preliminary versions before the publication date of the final version. Such advance electronic access does not advance the date of publication of a work, as preliminary versions are not published (Article 9.9)".

Gene Sequence Analysis of Mitochondrial COI in Genus *Saperda* (Coleoptera:Cerambycidae:Lamiinae)

Xu Wei^{1,2} Gan Qinhua³ Pu Jian⁴ Pan Yingwen² Cai Bo² Ao Su²

Meng Rui^{2*} Chen Li^{1*} Liu Fuxiu² Han Yuchun² Li Jiatong²

1.College of Plant Protection,Southwest University,Chongqing 400716,Chongqing municipality, China;2.Post-Entry Quarantine Station for Tropical Plants of Haikou Customs, Haikou 570311, Hainan Province, China; 3.Hainan University,Haikou 570228,Hainan Province,China; 4.HNPOLY PHARM. CO.,LTD.,Haikou 571127,Hainan Province,China)
The first author's E-mail:Xu Wei (840252833@qq.com)

Corresponding author E-mails:Meng Rui(541946422@qq.com) ;Chen Li (lichen57@swu.edu.cn)

Co-authors' E-mails:Gan Qinhua(32779939@qq.com);Pu Jian(312127516@qq.com);PanYingwen(71838341@qq.com);Cai Bo(81054870@qq.com);Ao Su (784808372@qq.com);Liu Fuxiu(58532901@qq.com);Han Yuchun(43016022@qq.com);Li Jiatong(372463464@qq.com)

Abstract: In this study, the total DNA of nine species of *Saperda* (*Lopezcolonia*) *octopunctata* (Scopoli, 1772), *Saperda* (*Lopezcolonia*) *scalaris* (Linnaeus, 1758), *Saperda* *interrupta* Gebler, *Saperda* *Alberti* (Plavilstshikov), *Saperda* (*Saperda*) *similis* Laicharting, 1784, *Saperda* (*Compsidia*) *populnea* (Linnaeus, 1758),*Saperda* (*Saperda*) *carcharias* (Linnaeus, 1758), *Saperda* (*Lopezcolonia*) *perforata* Pallas, 1773 and *Saperda ohbayashi* were extracted . Two partial sequences of mitochondrial gene and one partial sequence of nuclear gene were amplified. Comparing the COI sequence with the DNA barcode data in GenBank can effectively identify the related species of *Saperda*. It will be applied to the rapid identification of some species of *Saperda* in imported wood at ports, and improve the detection rate of plant quarantine.

Key words:Genus *Saperda*;mtDNA;COI;sequence analysis

Introduction

Saperda Fabricius, 1775, belongs to Insecta, Coleoptera, Cerambycidae, Lamiinae. There are more than 70 species of *Saperda* in the world and 23 species in China (Song and Chen 2008). For many years, many species of *Saperda* pests were often found in imported wood at ports. From October 2013 to October 2015, dozens of quarantine pests of *Saperda* (non Chinese species) were intercepted at ports. It is an urgent task to exclude the Chinese species of *Saperda* and intercept the quarantine pests of *Saperda* (non Chinese species). It is difficult and inaccurate for port inspection and quarantine personnel to exclude Chinese species of *Saperda* only from the external morphology, which often needs expert identification and review. The study

on the genetic differentiation and phylogeny of *Saperda* population can provide scientific basis for the study on the relationship between the source of *Saperda*, the occurrence mechanism and the accurate quarantine and identification of *Saperda*. It also has important theoretical and practical significance for improving the level of forecasting and monitoring technology of *Saperda* and formulating sustainable population control strategy.

Saperda (Lopezcolonia) octopunctata (Scopoli, 1772) is a non Chinese Quarantine Pest prohibited entry by the state. It was first intercepted from *Tilia* saw blade imported from Romania at Hainan port in 2008 and the male genitalia of seven species belonging to Genus *Saperda* was analyzed (Xu et al. 2008,2019). *Saperda ohbayashi* is an experimental specimen from Japan, but not distributed in China. It has been found in northern China that *Saperda (Lopezcolonia) scalaris* (Linnaeus, 1758)、*Saperda interrupta* Gebler、*Saperda alberti* (Plavilstshikov)、*Saperda (Saperda) similis* Laicharting, 1784、*Saperda (Compsidia) populnea* (Linnaeus, 1758)、*Saperda (Saperda) carcharias* (Linnaeus, 1758)、*Saperda (Lopezcolonia) perforata* Pallas, 1773 have been distributed. In this paper, the total DNA of these nine species of *Saperda* were extracted and the partial sequences of two mitochondrial genes and one nuclear gene were amplified. The COI sequences were compared with the DNA barcodes in GenBank to identify different species of *Saperda*.

1 Materials and Methods

1.1 Materials

Nine species specimens of *Saperda (Lopezcolonia) octopunctata* (Scopoli, 1772)、*Saperda (Lopezcolonia) scalaris* (Linnaeus, 1758)、*Saperda interrupta* Gebler、*Saperda alberti* (Plavilstshikov)、*Saperda (Saperda) similis* Laicharting, 1784、*Saperda (Compsidia) populnea* (Linnaeus, 1758)、*Saperda (Saperda) carcharias* (Linnaeus, 1758)、*Saperda (Lopezcolonia) perforata* Pallas, 1773 and *Saperda ohbayashi* were bought or presented from the cerambycids research experts who all collected the fresh specimens in 2015. If the samples received are not immersed in absolute ethanol, they should be immediately immersed in absolute ethanol and stored in - 80 °C refrigerator. The specimen information is shown in Table 1:

Table 1 Source information of *Saperda* specimens

Name specimen	of Host	Collecting place and time	Collector	Corresponding sequencing number
<i>Saperda octopunctata</i>	<i>Tilia</i>		J. Borucky	9
Scopoli		Czech Republic ,2015		
<i>Saperda alberti</i>	<i>Populus nigra</i>	Saima Town, Zang Kai		8
Plavilstshikov	var. <i>italia</i>	Fengcheng, Dandong, Liaoning, China		
<i>Saperda obhayashii</i>	Absent	2015-VI-14	Yamanashi	7
(Podany,1963)		Japan		
<i>Saperda interrupta</i>	Absent	2015-VI-11	Yamanashi	6
(Gepler,1825)		Japan		
<i>Saperda similis</i>	Larval rearing	2012-V-10	MORAVIA ex1	5
Laicharting			Bruntal-Razova	
			V. Sima	
<i>Saperda populea</i>	Absent	BOHEMIA		4
(Linnaeus)		Tatenice		
		2015-VI-11Czec		
		h Republic ,2015		
<i>Saperda perforata</i> Pallas	Larval rearing	BOHEMIA		3
		Czech		
		Republic ,2015		

<i>Saperda scalaris</i>	Larval rearing (Linnaeus)	BOHEMIA Czech Republic ,2015	Lanskrouneny	2
<i>Saperda carcharias</i>	<i>Salix caprea</i> (Linnaeus)	2015- V -18 Czech Republic ,2015	Vladimir Sima	1

1.2 Methods

1.2.1 Design and synthesis of primers

The universal primers designed by Simon et al. (1994) were used to amplify COI gene,C1-J-2183 (5'-CAACATTATTTGATTTTGG -3') and TL2-N-3014 (5'- TCCAATGCACTAATCTGCCATATTA -3') .

1.2.2 Extraction of genomic DNA

In this experiment, the leg or chest muscles of *Saperda* adult were used as experimental materials, and the phenol chloroform isoamyl alcohol method was used to extract DNA. The specific experimental steps were modified according to the extraction method of Sambrook et al. (1989).

1.2.3 PCR amplification and sequencing

PCR reaction system: 10 × PCR buffer (containing Mg2+) 2.5 μ L, 2.5mmol/l dNTP 2 μ L, Taq DNA polymerase (5U) 0.2 μ L, C1-J-2183/ TL2-N-3014 (10 μmol/L) 1 μ L, DNA template 1 μ L, water supplement to 25 μ L. The PCR reaction conditions were as follows: pre denaturation at 95 °C for 5 min, followed by 35 cycles: denaturation at 95 °C for 30 sec, annealing at 47 °C for 40 sec, extension at 72 °C for 1 min, extension at 72 °C for 5 min, and heat preservation at 4 °C. The amplified products were detected by 1% agarose gel electrophoresis and purified to Shenzhen Huagi Gene Technology Service Co., Ltd.

1.2.4 Sequence analysis

BioEdit v7.0.5, GeneDoC, Clustal W and MEGA 5.10 software were used to compare and analyze the sequences, and the phylogenetic tree was constructed.

2 Results and Analysis

2.1 Detection of PCR products

By agarose gel electrophoresis, 9 species of *Saperda* were amplified with clear and uniform bands (Fig.1), and negative control and blank control did not amplify strips. The results showed that the COI gene primer

C1-J-2183/ TL2-N-3014 could be used to amplify the gene fragments of 9 species of Cerambycidae, and the amplified fragments were consistent with the target fragments.




Fig.1 Agarose gel electrophoresis of PCR products amplified

2.2 Sequence composition and variation of different species of *Saperda*

The results showed that the average length of mtDNA CO I gene sequences of 9 different species of *Saperda* was 729 bases, the average contents of A, T, C and G were 30.9%, 38.1%, 16.1% and 14.9% respectively, the average contents of A + T and G + C were 69.0% and 31.0% respectively, and the average contents of A + T were more than 2 times of the average contents of G + C; the average contents of A + T at codon 1 were 62.6% and the average contents of A + T at codon 2 were 62.6%. The average content of C and G in locus 2 was 2.0% and 10.2%, respectively. The average content of A was the highest, reaching 44.1%, and the average content of T was 43.7%. The results showed that the frequency of base usage of codon was obviously biased. The contents of mtDNA co I a, t, C and G in different species of *Saperda* are different, as shown in the following table (Table 2). The results showed that there were significant genetic differences in base content among different species of *Saperda*.

Table 2 Comparison of base composition of COI gene in mtDNA of 9 species of *Saperda*

Species of <i>Saperda</i>	Sequence length/bp	base content /%			
		T	A	C	G
<i>Saperda octopunctata</i>	729	38.4	31.8	15.5	14.3
<i>Saperda alberti</i>	729	37.3	32.4	15.5	14.8
<i>Saperda ohbayashii</i>	729	38.4	28.5	17.3	15.8
<i>Saperda interrupta</i>	729	40.0	28.3	16.3	15.4
<i>Saperda similis</i>	729	38.3	31.3	16.0	14.4
<i>Saperda populnea</i>	729	36.8	31.8	17.0	14.4

<i>Saperda perforata</i> Pallas	729	37.7	31.3	15.5	15.5
<i>Saperda scalaris</i>	729	37.6	31.3	16.0	15.1
<i>Saperda carcharias</i>	729	38.1	31.3	16.2	14.4

There were 476 conserved sites and 253 variant sites in sequence length of 729 bp , with a variation rate of 34.71%. There were 171 parsimony informative loci and 82 self descendant loci, accounting for 32.41% of the total. The composition of gene base substitution of mtDNA CO I of nine different species of *Saperda* is shown in Table 3. Generally, the conversion between sequences is equivalent to transversion. Statistics show that the nucleotide substitution at different codon sites is different, especially at codon 2, the frequency of conversion and transversion is high. The conversion accounts for 61.82% of the total conversion, and the transversion accounts for 87.93% of the total transversion. It is a synonymous substitution, which causes less substitution of amino acids. The results show that the transversion mainly occurs between A and T, and there is almost no transversion between G and C; the transformation mainly occurs between T and C; the average ratio of transformation and transversion is 1.0. The frequency of transition and transversion of codon 1 is almost the same, and the frequency of transition of codon 3 is much higher than that of transversion.

Table 3 base substitution composition of mtDNA CO I gene in 9 species of *Saperda*

codon	ii	si	sv	R	Identical Pairs				Transitional Pairs			Transversional Pairs			Total	
					TT	CC	AA	GG	TC/CT	AG/GA	TA/A	TG/GT	CA/AC	CG/GC		
1	238	2	2	1.1	104	52	46	37	1/1	0/0	0/0	0/0	0/0	0/0	243	
2	158	34	51	0.7	72	6	80	0	13/14	3/3	21/19	1/1	4/5	0/0	243	
3	220	19	5	3.9	58	33	66	63	7/6	3/3	2/1	0/0	0/1	0/0	243	
Averag	615	55	58	1.0	234	91	191	100	21/21	7/7	23/20	1/1	5/6	1/1	729	
e																

Notes: ii (Identical Pairs) ; si (Transitional Pairs) ; sv (Transversional Pairs) ; R=Transitional Pairs/Transversional Pairs

2.3 Analysis of genetic distance among different species of *Saperda*

Based on the Kinura-2-Parameter model, the genetic distances among different species of *Saperda* were analyzed (Table 4). The average genetic distance of the nine species was 0.271, while that of *Saperda similis* and *Saperda carcharias* was the smallest, only 0.001. The results showed that the genetic distance between *Saperda interrupta* and *Saperda alberti* was the largest (0.927), the genetic distance between *Saperda interrupta* and *Saperda populnea* was the second largest (0.909), and the genetic distance between *Saperda*

octopunctata and *Saperda interrupta* and that between *Saperda octopunctata* and *Saperda ohbayashii* was 0.631 and 0.594 respectively, which were a little larger.

Table 4 Genetic distance of mtDNA COI gene sequences of 9 species of *Saperda*

Species	of	9	8	7	6	5	4	3	2	1
<i>Saperda</i>										
9										
8		0.066								
7		0.594	0.731							
6		0.631	0.927	0.141						
5		0.023	0.095	0.399	0.484					
4		0.182	0.111	0.565	0.909	0.128				
3		0.084	0.067	0.410	0.556	0.066	0.171			
2		0.071	0.060	0.372	0.560	0.034	0.086	0.018		
1		0.032	0.092	0.390	0.501	0.001	0.104	0.067	0.029	

Notes: 1.*Saperda carcharias*; 2.*Saperda scalaris*; 3.*Saperda perforata*; 4.*Saperda populnea*; 5.*Saperda similis*; 6.*Saperda interrupta*; 7.*Saperda ohbayashi*; 8.*Saperda alberti*; 9.*Saperda octopunctata*

2.4 Construction of phylogenetic analysis

One sequence of 9 species of longicorn was selected and analyzed by mega4.0 software. The neighbor joining method (N-J method), maximum parsimony method (M-P method) and UPGMA method were used to draw the diagrams respectively. The confidence of each branch was tested by "bootstrap" 1000 times. The obtained tree diagrams are respectively Fig. 2, Fig. 3 and Fig. 4. From the topological structure, the results of the three clustering methods are basically consistent. The results showed that nine species of *Saperda* were clustered into one branch, and the same species were all independent, and the self expanding support rate was 100%. The similarity between the two species *Saperda similis* and *Saperda carcharias* was the highest, which was aggregated with 100% confidence, indicating that the two species were closely related. In the phylogenetic tree, *Saperda scalaris* and *Saperda interrupta* were clustered into one branch with high confidence, indicating that they were closely related. There are obvious differences among different species of *Saperda*, indicating that this sequence can be used as the basis for molecular identification of *Saperda*.




Fig. 2 N-J tree based on the analysis of COI gene sequence of the 9 *Saperda* species





Fig. 3 M-P tree based on the analysis of COI gene sequence of the 9 *Saperda* species



0.10 0.08 0.06 0.04 0.02 0.00

Fig. 4 UPGMA tree based on the analysis of COI gene sequence of the 9 *Saperda* species

3 Discussion

3.1 DNA barcoding can be used to identify the species of *Saperda*

DNA barcoding technology needs two conditions to identify species: first, each species has a unique DNA barcode and there is no shared sequence among different species; second, the genetic distance between species should be much greater than that within species. In this study, the COI sequences of 9 species of *Saperda* were amplified by universal primers, and a fragment of 729bp was obtained. The results show that the genetic distance between species was significantly greater than that within species.

There was no overlap between intraspecific genetic distance and interspecific genetic distance, and interspecific genetic distance was significantly greater than intraspecific genetic distance. Based on the DNA barcode sequence, the phylogenetic tree was constructed by using the adjacency method and the maximum parsimony method. The results showed that the same species clustered into a small branch, and the node support rate was 100%. The results of genetic distance and phylogenetic tree showed that the fragment could be used as the basis for the analysis and identification of Cerambycidae species.

3.2 Validation of classification results using molecular barcode data

There are many species of *Saperda* with a long history of classification. Due to intraspecific variation, such as color and stripe variation, it is often difficult to accurately and quickly identify the species of *Saperda*. The results of NJ tree and MP tree showed that a species of *Saperda* was an independent species at the molecular level.

Due to the limitation of specimen, only 9 species of *Saperda* were studied in this study. There are still many problems to be clarified in the classical taxonomy of *Saperda* based on external morphology and male genitalia. The external morphological characteristics of some species are very similar. For example, the color and spot position of *Saperda octopunctata*, *Saperda punctata*, *Saperda perforata* Pallas and *Saperda alberti* Plavilstshikov species are easy to confuse. In the future, molecular barcode data should be supplemented on the basis of morphological classification and accurate identification, and NCBI and bold database should be used for large-scale research in order to identify and analyze the phylogenetic relationship of Cerambycidae more quickly and accurately (Kethidi D R et al. 2003; Ma Jian et al. 2010; Li Jing et al. 2011, 2013; Wei Zihan et al. 2014; Zheng Sizhu et al. 2015; Zhang Shuang et al. 2015)

Acknowledgements

We thank Li Zhu (College of Plant Protection of Southwest University) for her careful reading of our manuscript and her insightful suggestions. This project is supported by the funds of Science Technology Project (2014IK015, 2017IK203) of the former General Administration of State Quality Supervision and Inspection and Quarantine and Hainan Natural Science Fund Youth Project Fund (319QN334).

References

- Kethidi DR, Roden DB, Ladd TR, et al. (2003) Development of SCAR markers for the DNA-based detection of the Asian long-horned beetle, *Anoplophora glabripennis* (Motschulsky). Archives of Insect Biochemistry and Physiology 52(4): 193-204.
- Li J, An SH, Yin XM (2011) Phylogeny of 12 species of Lamiinae beetles based on COI gene sequence. Vol. 7. Central China insect research, 243.
- Li J, An SH, Zhang HF, et al. (2013) Characteristics and phylogeny of mitochondrial Co II gene in some species of Lamiinae beetles. Journal of Henan Agricultural University 47 (6): 715-721.
- Ma J, Li CD, Deng LW, et al. (2010) Assisted identification of several poplar stem borer larvae based on COI gene restriction endonuclease. Journal of Northeast Forestry University 38 (5): 83-87.
- Sambrook J, Fritsch EF and Maniatis T. Molecular Cloning: A Laboratory Manual. Cold Spring Harbor Press, Cold Spring Harbor, New York.
- Simon M, Bénit P, Vassal A, Dubois C, Faye G (1994) Sequence of the PHO2-POL3 (CDC2) region of chromosome IV of *Saccharomyces cerevisiae*. Yeast 10(12):1653-6.
- Song YQ, Chen L. A Preliminary Fauna Analysis of Saperdini in China (Coleoptera: Cerambycidae: Lamiinae). Southwest China Journal of Agricultural Sciences, 2008, 21(2):498-502.
- Xu W, Cai B, Ao S, Liu FX, Pan YW, Han YC, Lin MG, Chen L (2019) A Comparative Study on the Characters of the Male Genitalia of 7 Species of *Saperda* spp. Journal of Environmental Entomology 41 (3):581-591.
- Xu W, Lin MG, Han YC, Nian XL (2008) The First Quarantine Pest, *Saperda octopunctata*, Intercepted in China. Plant Quarantine 22(6):384.
- Wei ZH, Yin XM, An SH, Su LJ, Li J, Zhang HF (2014) Molecular phylogenetic study of the higher taxa of the superfamily Cerambycoidea (Insecta: Coleoptera) based on the combined analysis of ribosomal DNA sequences. Acta Entomologica Sinica 57(6):710-720.
- Zheng SZ, Zhang K, Yang XJ, et al. (2015) Comparison of COI genes and preliminary systematic study on Monochamus. Acta Entomologica Sinica 52 (2): 370-381.

Zhang S, Su XY, Huang DZ, et al. Sequence variation and genetic differentiation of mitochondrial COI gene in different geographic populations of *Germari* cerambycids. Sericultural Science, 2015, 41 (2): 234-238