

## PREPRINT

*Author-formatted, not peer-reviewed document posted on 29/11/2022*

DOI: <https://doi.org/10.3897/arphapreprints.e98091>

# Species associated with whale sharks (*Rhincodon typus*) in the Galapagos Archipelago

Sofia Green, Alex Hearn, Jonathan Green

### Disclaimer on biological nomenclature and use of preprints

The preprints are preliminary versions of works accessible electronically in advance of publication of the final version. They are not issued for purposes of botanical, mycological or zoological nomenclature and **are not effectively/validly published in the meaning of the Codes**. Therefore, nomenclatural novelties (new names) or other nomenclatural acts (designations of type, choices of priority between names, choices between orthographic variants, or choices of gender of names) **should NOT be posted in preprints**. The following provisions in the Codes of Nomenclature define their status:

### International Code of Nomenclature for algae, fungi, and plants (ICNafp)

**Article 30.2:** "An electronic publication is not effectively published if there is evidence within or associated with the publication that its content is merely preliminary and was, or is to be, replaced by content that the publisher considers final, in which case only the version with that final content is effectively published." In order to be validly published, a nomenclatural novelty must be effectively published (Art. 32.1(a)); in order to take effect, other nomenclatural acts must be effectively published (Art. 7.10, 11.5, 53.5, 61.3, and 62.3).

### International Code of Zoological Nomenclature (ICZN)

**Article: 21.8.3:** "Some works are accessible online in preliminary versions before the publication date of the final version. Such advance electronic access does not advance the date of publication of a work, as preliminary versions are not published (Article 9.9)".

# Species associated with whale sharks (*Rhincodon typus*) in the Galapagos Archipelago

Sofia M Green<sup>‡</sup>, Alex Hearn<sup>§,‡</sup>, Jonathan R Green<sup>‡</sup>

<sup>‡</sup> Galapagos Whale Shark Project, Puerto Ayora, Ecuador

<sup>§</sup> Universidad San Francisco de Quito, Quito, Ecuador

Corresponding author: Sofia M Green ([sgreenitu@gmail.com](mailto:sgreenitu@gmail.com))

## Abstract

Whale sharks (*Rhincodon typus*) frequently appear to interact or associate with other species, which vary depending on the community structure and the demographic of the whale sharks at each location globally. Here, we present the species sighted frequently around whale sharks in the Galapagos Archipelago, and reported by dive guides and scientists and also in earlier publications. These associated species include cetacean species: bottlenose dolphins (*Tursiops truncatus*), other shark species: silky sharks (*Carcharhinus falciformis*), Galapagos sharks (*Carcharhinus galapagensis*), scalloped hammerhead sharks (*Sphyrna lewini*), tiger sharks (*Galeocerdo cuvier*), and teleost fish species: remoras (*Echeneis sp*), yellowfin tuna (*Thunnus albacares*), almaco jacks (*Seriola rivoliana*), and black jacks (*Caranx lugubris*). The recording of interspecies associations and interactions may lead to better understanding of the natural history of whale sharks and can show important symbiotic relationships or interdependence between different species.

## Keywords

Species associations, interspecific interactions, natural history, symbiosis, parasitism

## Introduction

The whale shark, *Rhincodon typus* (Smith, 1828), is a circumglobal tropical and temperate water species (Rowat and Brooks 2012). It is mostly solitary and pelagic, although it also forms seasonal feeding aggregations of mostly immature males in at least a dozen (mostly coastal) sites (e.g. Belize, Heyman et al. 2001; Gulf of Mexico, Hoffmayer et al. 2007; Tanzania, Rohner et al. 2015). Several species are known to associate with whale sharks around the globe, yet they vary across different locations. Bottlenose dolphins, *Tursiops truncatus*, have been reported swimming with whale sharks or feeding alongside them in the Gulf of Mexico, Belize, Honduras, the Red Sea and Christmas Island (Latusek-

Nabholz 2014). Cleaning interactions have been reported by Araujo et al. (2020) of the blue-streak cleaner wrasse, *Labroides dimidiatus*, and the moon wrasse, *Thalassoma lunare*, in the Philippines and by Quimbayo et al. (2016) of the King Angelfish, *Holocanthus passer*, in Malpelo Island. Other associations recorded include suckerfish (*Remora* sp. and *Echeneis nucrates*) in North-Western Australia (Norman 1999), schooling fish such as tuna and mackerels, plus carcharhinid shark species swimming alongside whale sharks off South Texas (Hoffman et al. 1981) and more tuna species associations in the Atlantic and Indian Oceans (Escalle et al. 2018). Indeed, until recently, purse seine vessels would set their nets around whale sharks, in the knowledge that there would likely be tuna schooling around them. The use of whale sharks as living Fish Aggregation Devices (FADs) is now banned by most tuna regional fisheries management organizations (RFMOs), including in the Eastern Pacific (Resolution C-19-06, IATTC 2019) where it is prohibited to set a purse-seine net around a school of tuna known to be swimming around a living whale shark.

The Eastern Pacific hosts the only known whale shark aggregation made up almost exclusively of large females – the Galapagos Islands (Norman et al. 2017). The first report of a whale shark in the Galapagos was recorded by William Beebe in 1925 of an individual swimming under the stern of a boat on the Arcturus expedition (reported in Gudger 1933). They continued to be reported occasionally by guides between Floreana and Isabela, in the Canal Bolivar, and around other few spots around the Archipelago (see Fig. 1). Prior to whale shark focused tourism, a short note was written of species associated with whale sharks in the Galapagos Islands by a group of cetacean scientists who spotted four whale sharks by chance during their tracking of sperm whales off the coast of southwest Isabela in 1988. They reported sighting these whale sharks with remoras, *Echeneis* sp., around and in the sharks mouth, hammerhead sharks, *Sphyrna lewini*, and tiger sharks, *Galeocerdo cuvier*, swimming close by (at 10 m distance) a whale shark, and yellowfin tuna, *Neothunnus macropterus/ Thunnus albacares*, surrounding a whale shark (Arnbom and Papastavrou 1988).

Although seen sporadically throughout the archipelago, whale sharks occur predictably at the northernmost island of Darwin, especially in the months of July through November (Acuña-Marrero et al. 2014), and have become an important dive tourism attraction for live-aboard vessels during these months. Since 2011, a team of scientists have led yearly expeditions to Darwin Island to record whale shark movements, behaviour, and to attempt to establish their reproductive state. The team also began to collect sightings reports from dive guides from this time. Here, we report on the association of different species with whale sharks at Darwin and across the Galapagos Marine Reserve, in the Eastern Tropical Pacific.

## Methods

## Study Site

The Galapagos Archipelago and its Marine Reserve are a UNESCO Natural World Heritage Site, and possess a unique set of characteristics which makes it a hotspot for biodiversity which must be protected. Its isolation from the continent, lying approximately 1000 km due west of mainland Ecuador, and its location straddling the equator, allows for species diversification, which has led to its unique wildlife, with 18% of marine species being endemic (Bustamante et al. 2000). The marine currents that influence the islands are also important for the diversity of life on the islands. The Panama current brings warmer waters and carries species from the north, the Humboldt Current brings colder waters and species from the Antarctic and sub polar region, and the Equatorial Under-Current, (also known as the Cromwell Current) brings cold, nutrient rich water, which upwells along the western margins of the archipelago. This upwelling generates one of the most productive marine ecosystems in the world (Schaeffer et al. 2008, Smith and Khaled bin Sultan Living Oceans Foundation 2012), which in turn attracts highly migratory species such as the whale shark.

Darwin and Wolf Islands are in the far northern region of the archipelago, and predictable seasonal sightings of whale sharks occur between July-December of every year (Acuña-Marrero et al. 2014).

## Data Collection

For the past decade (2011-2021) the Galapagos Whale Shark Project has led yearly expeditions to this area and collected data on the physiology and ecology of whale sharks and has noted species interactions used for this study. The data collected by the project team come from performing three dives a day every day for two weeks every year.

Data from dive guides and fishermen was collected informally throughout the years. All dive liveboards in the Galapagos have both Darwin and Wolf in their itineraries where dive guides spend 1-2 days, with 3-4 dives per day, every week in these waters. The dive guides send sporadic informal reports on some notable encounters they have with whale sharks and other species. Fishermen data on the other hand comes mostly from the southern and central islands of the reserve where most of their fishing activities occur. The data collected from their sightings comes from word of mouth accounts.

All data presented in this report were collected between 2010-2020, except the associations recorded by Arnbom and Papastavrou (1988), which we also include in this note.

During this study there is mention of species interactions and species associations. For the purposes of this report, we defined associations as an encounter of a species other than *Rhincodon typus*, occurring within 10 m of a whale shark and moving in the same direction, or physically interacting with the shark. Meanwhile, in ecology, species interactions have been defined as the relationship between two or more species that form an

interdependence to some degree. The five major types of species interactions have been labelled as predation, competition, mutualism, commensalism and amensalism (Hillis et al. 2011). Thus, species interactions would be classified as a species association, yet not all associations would be defined as species interactions.

## Results

Nine species, including four other shark species, were found in association with whale sharks in the Galapagos Islands (Table 1) as reported by the GWSP team, dive guides and fishermen of the reserve.

Whale sharks are frequently sighted with other shark species swimming alongside such as *Sphyrna lewini*, *Carcharhinus galapagensis*, *Carcharhinus limbatus*, and *Carcharhinus falciformis*. However, *C. galapagensis* and *C. falciformis* have both displayed unusual behaviour around whale sharks not reported elsewhere. Both *Carcharhinus* sp. have been sighted rubbing themselves on the whale sharks' body and heads on multiple occasions and reported by several Galapagos National Park Guides and the GWSP scientists. This is supported with photographic and video evidence as seen in figure 2. Both shark species are thought to do this for cleaning purposes, to rid themselves of ectoparasites. Recently, *C. galapagensis*, particularly females, have also been sighted trailing close behind whale shark tails in large groups (4+ individuals, refer to Fig. 2) and displaying agonistic behaviour towards divers when approaching the whale shark (Sofia M Green, pers. observation).

Associations between dolphins and whale sharks have been reported previously by Latusek-Nabholz (2014), off the Gulf of Mexico and have been noted again in the Galapagos Archipelago. Bottlenose dolphins, *Tursiops truncatus*, are sighted almost every day at Darwin Island, leading scientists to believe there may be a resident population (data deficient). These dolphins are frequently sighted during dives at Darwin Arch.

The GWSP has recorded on a number of occasions, particularly in September 2018 and August 2020, dolphins swimming alongside the whale shark and seemingly bow riding its head (Fig. 3). Dive guide Paulo Tobar reported the same behaviour during once of his expeditions to Darwin in 2018.

Remora or suckerfish associations have also been reported previously associated with whale sharks around the globe (Cohen et al. 2020, Norman 1999, Colman 1997). This association has previously been noted in the Galapagos Archipelago by Arnbom & Papastavrou in 1988 and again by Galapagos National Park guides and scientists from the GWSP. Remora's, *Echeneis* sp., are seen on almost all whale sharks sighted in the islands. They are generally around and inside the whale shark's mouth, gills, and cloaca (see Fig. 4).

Two species of schooling fish have also been sighted frequently in association with the whale shark. In the Galapagos Islands, whale sharks are rarely seen without the company of the Black Jacks, *Caranx lugubris*, and also frequently sighted with the Almaco Jack,

*Seriola rivoliana*. These fish are usually surrounding the shark, and swimming close by to its body (Fig. 4).

## Discussion

Associations between different marine species arise naturally from sharing an environment. In this study we identified nine species which associate with whale sharks in the Galapagos Marine Reserve. Only one of these appeared to be a long-term association – the remoras affixed to the bodies of the sharks. Two shark species interacted physically with the whale sharks, apparently using their abrasive skin as cleaning tools. The remainder of the associations did not involve physical contact with the shark, but either involved swimming close to the body of the shark (as with the jacks and tunas) or utilizing disruptions in water movements caused by the sharks' movements, as with the dolphins bow riding or the silky sharks swimming in the slipstream. The agonistic display towards the diver by the Galapagos shark may hint to a more meaningful significance for this association with whale sharks. The agonistic behaviour, caused by the rapidly approaching divers was equivalent to the behaviour initially described by Johnson and Nelson (1973) for grey reef sharks and later reinforced by Martin (2007) as noted on other *Carcharhinus* species.

As mentioned by Rowat and Brooks (2012) and reported in this study the sucker fish species not only attach to the whale shark but live in their mouth, gill slits, cloaca, peribranchial cavity and the spiracle. Although there has been some discussion of whether this symbiosis is mutualistic, parasitic or commensalistic, we report it as parasitic, defined as the relationship where one individual benefits at the expense of the other organism, from sightings observed in the Galápagos. The suckerfish have been observed removing parts from inside the gills, covering the entire mouth, and entering the reproductive areas of the sharks, most likely causing damage to the whale shark.

The behaviour of the bottlenose dolphins in the Galapagos, where they are sighted bow riding the shark, is similar as the behaviour reported in the Gulf of Mexico, Belize, Honduras, the Red Sea and Christmas Island, where the mammals were sighted swimming alongside the whale shark (Latusek-Nabholz 2014).

The interactions in each sighting location is based on which species exist naturally in the area and the habitat use of the whale shark. For instance, earlier we mentioned the cleaning behaviour of reef fish, the blue-streak cleaner wrasse, *Labroides dimidiatus*, and the moon wrasse, *Thalassoma lunare*, recorded in the Philippines and of the King Angelfish, *Holocanthus passer*, in Malpelo Island (Araujo et al. 2020). In Galapagos, although both these species are present, we have not sighted the same behaviour from the cleaner fish with whale sharks, as we do with hammerhead sharks. We suggest this would most likely be due to the fact that the adult whale sharks in the Galapagos Archipelago are sighted in the open ocean, and do not spend a significant amount of time on or close to the reefs, while the juvenile whale sharks sighted in the Philippines would. Meanwhile, as mentioned by Araujo et al. (2020), the single sighting of cleaning behaviour in Malpelo

Island by the King Angelfish, where whale sharks are also known to be transient and pelagic, is rare and probably opportunistic.

## Acknowledgements

We would like to thank the naturalist and dive guides who send in reports with a special thank you to Paulo Tobar for always contributing information to the project and incentivising his co-workers to do so as well. We would also like to thank the fishermen Manuel Yopez and Daniel Torres who have sent information about sighting locations and interesting encounters and have kept us informed about the sightings reported by their colleagues.

## Ethics and security

All data collection is conducted under, and approved by, the Galapagos National Park Permits under the project “The role of oceanic islets for the protection of marine migratory species”.

## Conflicts of interest

No conflicts of interest.

## References

- Acuña-Marrero D, Jiménez J, Smith F, Doherty P, Hearn A, Green J, Paredes-Jarrín J, Salinas-de-León P (2014) Whale Shark (*Rhincodon typus*) seasonal presence, residence time and habitat use at Darwin Island, Galapagos Marine Reserve. PLOS One 9 (12). <https://doi.org/10.1371/journal.pone.0115946>
- Araujo G, Miranda J, Allen H, Labaja J, Snow S, Ponzó A, Legaspi C (2020) Whale sharks *Rhincodon typus* get cleaned by the bluestreak cleaner wrasse *Labroides dimidiatus* and the moon wrasse *Thalassoma lunare* in the Philippines. Journal of Fish Biology 97 (4): 1247-1251. <https://doi.org/10.1111/jfb.14464>
- Arnbom T, Papastavrou V (1988) Fish in association with whale sharks *Rhincodon typus* near the Galapagos Islands. Noticias de Galapagos 46: 13-15.
- Bustamante R, Collins KJ, Bensted-Smith R (2000) Biodiversity Conservation in the Galapagos Marine Reserve. Bulletin De L'institut Royal Des Sciences Naturelles De Belgique 70: 31-38.
- Cohen K, Flammang B, Crawford C, Hernandez LP (2020) Knowing when to stick: touch receptors found in the remora adhesive disc. Royal Society Open Science 7 (1). <https://doi.org/10.1098/rsos.190990>
- Colman JG (1997) A review of the biology and ecology of the whale shark. Journal of Fish Biology 51 (6): 1219-1234. <https://doi.org/10.1111/j.1095-8649.1997.tb01138.x>
- Escalle L, Gaertner D, Chavance P, Murua H, Simier M, Pascual-Alayón PJ, Ménard F, Ruiz J, Abascal F, Mérigot B (2018) Catch and bycatch captured by tropical tuna purse-

- seine fishery in whale and whale shark associated sets: comparison with free school and FAD sets. *Biodiversity and Conservation* 28 (2): 467-499. <https://doi.org/10.1007/s10531-018-1672-1>
- Gudger EW (1933) A second whale shark, *Rhincodon typus*, at the Galapagos Islands. *Nature* 132 (3336): 569-569. <https://doi.org/10.1038/132569a0>
  - Heyman W, Graham R, Kjerfve B, Johannes R (2001) Whale sharks *Rhincodon typus* aggregate to feed on fish spawn in Belize. *Marine Ecology Progress Series* 215: 275-282. <https://doi.org/10.3354/meps215275>
  - Hillis DM, Heller HC, Price MV (2011) *Principles of Life*. Macmillan Learning
  - Hoffman W, Fritts T, Reynolds R (1981) Whale sharks associated with fish schools off South Texas. *Northeast Gulf Science* 5 (1). <https://doi.org/10.18785/negs.0501.08>
  - Hoffmayer E, Franks J, Driggers W, Oswald K, Quattro J (2007) Observations of a feeding aggregation of whale Sharks, *Rhincodon typus*, in the North Central Gulf of Mexico. *Gulf and Caribbean Research* 19 <https://doi.org/10.18785/gcr.1902.08>
  - IATTC (2019) Resolution C-19-06. Conservation of whale Sharks. [https://www.iattc.org/PDFFiles/Resolutions/IATTC/English/C-19-06-Active Whale%20sharks.pdf](https://www.iattc.org/PDFFiles/Resolutions/IATTC/English/C-19-06-Active%20Whale%20sharks.pdf)
  - Johnson R, Nelson D (1973) Agonistic Display in the Gray Reef Shark, *Carcharhinus menisorrah*, and Its Relationship to Attacks on Man. *Copeia* 1973 (1). <https://doi.org/10.2307/1442360>
  - Latusek-Nabholz J (2014) A rare observation of bottlenose dolphins (*Tursiops truncatus*) with a whale shark (*Rhincodon typus*) off the Florida Panhandle, Gulf of Mexico. *Aquatic Mammals* 40 (3): 293-296. <https://doi.org/10.1578/am.40.3.2014.293>
  - Martin RA (2007) A review of shark agonistic displays: comparison of display features and implications for shark–human interactions. *Marine and Freshwater Behaviour and Physiology* 40 (1): 3-34. <https://doi.org/10.1080/10236240601154872>
  - Norman B, Holmberg J, Arzoumanian Z, Reynolds S, Wilson R, Rob D, Pierce S, Gleiss A, de la Parra R, Galvan B, Ramirez-Macias D, Robinson D, Fox S, Graham R, Rowat D, Potenski M, Levine M, Mckinney J, Hoffmayer E, Dove AM, Hueter R, Ponzo A, Araujo G, Aca E, David D, Rees R, Duncan A, Rohner C, Prebble CM, Hearn A, Acuna D, Berumen M, Vázquez A, Green J, Bach S, Schmidt J, Beatty S, Morgan D (2017) Undersea Constellations: The global biology of an endangered marine megavertebate further informed through citizen science. *BioScience* 67 (12): 1029-1043. <https://doi.org/10.1093/biosci/bix127>
  - Norman BM (1999) Aspects of the biology and ecotourism industry of the whale shark *Rhincodon typus* in North-Western Australia. Masters by Research thesis, Murdoch University
  - QGIS Development Team (2020) QGIS Geographic Information System. Open Source Geospatial Foundation Project. URL: <http://qgis.osgeo.org>
  - Quimbayo JP, Dias MS, Schlickmann ORC, Mendes TC (2016) Fish cleaning interactions on a remote island in the Tropical Eastern Pacific. *Marine Biodiversity* 47 (2): 603-608. <https://doi.org/10.1007/s12526-016-0493-2>
  - Rohner C, Armstrong A, Pierce S, Prebble CM, Cagua EF, Cochran JM, Berumen M, Richardson A (2015) Whale sharks target dense prey patches of sergestid shrimp off Tanzania. *Journal of Plankton Research* 37 (2): 352-362. <https://doi.org/10.1093/plankt/fbv010>

- Rowat D, Brooks KS (2012) A review of the biology, fisheries and conservation of the whale shark *Rhincodon typus*. *Journal of Fish Biology* 80 (5): 1019-1056. <https://doi.org/10.1111/j.1095-8649.2012.03252.x>
- Schaeffer B, Morrison J, Kamykowski D, Feldman G, Xie L, Liu Y, Sweet W, McCulloch A, Banks S (2008) Phytoplankton biomass distribution and identification of productive habitats within the Galapagos Marine Reserve by MODIS, a surface acquisition system, and in-situ measurements. *Remote Sensing of Environment* 112 (6): 3044-3054. <https://doi.org/10.1016/j.rse.2008.03.005>
- Smith J, Khaled bin Sultan Living Oceans Foundation (2012) Galapagos Ocean Currents. <https://www.livingoceansfoundation.org/galapagos-currents/>






Figure 2.

Association of requiem sharks with whale sharks a) Image capture from BBC bodycam of a silky shark (*C. falciformis*) rubbing on a whale shark (*R. typus*) b) Image capture of Galapagos sharks (*C. galapagensis*) rubbing on a whale shark and swimming beside it (Photo credit GWSP)




Figure 3.

Bottlenose dolphins (*T. truncatus*) bow riding whale shark's head a)2018 b)2020 (Photo credit GWSP)




Figure 4.

Teleost fish species association a) Remoras (*Echeneis* sp.) inside the whale shark's gills, b) Remoras on mouth with both jack species swimming around, and c) Almaco jacks surrounding a whale shark (Photo credit GWSP)

Table 1.

Species associated with *R. typus* reported in the Galapagos Marine Reserve.

#	Species Associated		Report year	Reported by	# Reports	Behaviour
	Common Name	Scientific name				
1	Silky sharks	<i>Carcharhinus falciformis</i>	2016-2020	GWSP team, Galapagos dive guides	>10	Sighted frequently rubbing on whale shark's body and head.
2	Galapagos sharks	<i>Carcharhinus galapagensis</i>	2019, 2020	GWSP team, Dive guide: Paulo Tobar	>20	Observed following closely behind sharks. Occasionally also sighted rubbing on the body and head of the whale shark.
3	Hammerhead sharks	<i>Sphyrna lewini</i>	1988	Arnbom and Papastavrou 1988	100+	Observed swimming within 10 m of whale shark.
4	Tiger shark	<i>Galeocerdo cuvier</i>	1988	Arnbom and Papastavrou 1988	1	Observed swimming within 10 m of whale shark.
5	Bottlenose dolphins	<i>Tursiops truncatus</i>	2018-2020	Dive guide: Paulo Tobar, GWSP team	3	Sighted bow-riding whale sharks twice in 2018 and once in 2020. Frequently observed swimming around the shark.
6	Remora	<i>Echeneis</i> sp. posib. <i>naucrates</i>	1988, 2010-2020	Arnbom and Papastavrou 1988, GWSP team, Dive guide: Natalia Cifuentes	>100	Attached to whale sharks' body. Seen mostly around and inside the mouth, caudal fin, pelvic fins of whale sharks. Also, spotted inside the gill slits and the cloaca of adult females. Known to do so as an efficient form of transportation and to feed off of left-over food particles.
7	Black jacks	<i>Caranx lugubris</i>	2010-2020	GWSP team	>100	Swim alongside <i>R. typus</i> , possibly making use of slipstream.
8	Almaco jacks	<i>Seriola rivoliana</i>	2020	GWSP team and fishermen	>50	Swim alongside <i>R. typus</i> , possibly making use of slipstream.
9	Yellowfin tuna	<i>Neothunnus macropterus</i> / <i>Thunnus albacares</i>	1988	Arnbom and Papastavrou 1988	1	Surrounding <i>R. typus</i> .